EBV LMP1 in Gingival Epithelium Potentially Contributes to Human Chronic Periodontitis *via* Inducible IL8 Production

NORIHISA WATANABE^{1,2}, KEIKO NODOMI², RYO KOIKE^{2,3}, AYAKO KATO⁴,
OSAMU TAKEICHI⁵, AI KOTANI⁶, TADAYOSHI KANEKO³, HIROSHI SAKAGAMI⁷,
MASAMI TAKEI⁸, YORIMASA OGATA⁴, SHUICHI SATO¹ and KENICHI IMAI²

¹Department of Periodontology, Nihon University School of Dentistry, Tokyo, Japan;

²Department of Microbiology, Nihon University School of Dentistry, Tokyo, Japan;

³Department of Oral and Maxillofacial Surgery, Nihon University School of Dentistry, Tokyo, Japan;

⁴Department of Periodontology, Nihon University School of Dentistry at Matsudo, Chiba, Japan;

⁵Department of Endodontics, Nihon University School of Dentistry, Tokyo, Japan;

⁶Department of Hematological Malignancy, Institute of Medical Science, Tokai University, Kanagawa, Japan;

⁷Meikai University Research Institute of Odontology (M-RIO), Saitama, Japan;

⁸Division of Hematology and Rheumatology, Department of Medicine,
Nihon University School of Medicine, Tokyo, Japan

Abstract. *Background/Aim:* Human chronic periodontitis is a major health problem. Although some oral bacteria have been reported to be putative pathogens, Epstein–Barr virus (EBV) is reported to be associated with the progression of periodontitis. However, the role of EBV in the aetiology of periodontitis is unknown. Therefore, we investigated periodontal pathogenesis of EBV to confirm whether EBV-encoded latent membrane protein 1 (LMP1) induces Interleukin-8 (IL8) production in human gingival cells. *Materials and Methods:* Real-time polymerase chain reaction, luciferase assay, enzyme-linked immunosorbent assay (ELISA), and western blotting were performed for determining IL8 mRNA expression, nuclear factor kappa B (NF- κ B) transcription, IL8 production, and the phosphorylation of NF- κ B p65 and Inhibitor of kappa B alpha (I κ B α), respectively, in Ca9-22 human gingival epithelial cells. Two LMP1 mutants lacking C-terminal activating region (CATR) domains responsible for activating NF- κ B were used. *Results:* Extremely high IL8 production was induced by LMP1 in time- and dose-dependent manner, where simultaneous phosphorylation of NF- κ B p65 and I κ B α

and transcription of NF- κ B were observed. On the contrary, IL8 production and NF- κ B transcription were drastically inhibited by dominant negative mutant of I κ B α . Moreover, the LMP1 mutants failed to induce IL8 production. *Conclusion:* Our findings suggest that due to CATR domains, LMP1 contributes to the progression of periodontitis via IL8 production attributable to NF- κ B activation.

Chronic periodontitis, a chronic inflammatory and infectious disease causing the destruction of the periodontium including the alveolar bone, is prevalent worldwide (1, 2). Mounting evidence has indicated that chronic periodontitis is a risk factor for pre-term birth, heart disease, diabetes, and atherosclerosis (1, 2). Over the past decade, neutrophil infiltration in the periodontium has been revealed to be the major aetiology for periodontitis (2). Some oral endogenous bacteria are believed to trigger periodontitis *via* host–parasite interactions (2, 3). However, periodontopathic bacteria, such as *Porphyromonas gingivalis*, are not always detected in periodontal lesions (4–6); therefore, the conventional theory based on bacterial aetiology alone cannot fully explain the aetiology of periodontitis.

A positive association has been reported between chronic periodontitis and Epstein–Barr virus (EBV) infection (7–11). EBV, a member of the herpesvirus family, infects many adults. During primary EBV infection, the virus undergoes lytic replication in B-cells and epithelial cells of the upper aerodigestive tract, where it later establishes latency (12, 13). EBV can be reactivated and is commonly found in the saliva of infected people (9, 10, 14). Many reports have demonstrated that the amount of EBV DNA detected in

This article is freely accessible online.

Correspondence to: Kenichi Imai, 1-8-13 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-8310, Japan. Tel: +81 332198115; Fax: +81 332198317, e-mail: imai.kenichi@nihon-u.ac.jp

Key Words: EBV, LMP1, periodontitis, IL8, NF- κ B, gingival epithelial cells.

periodontal pockets of patients with chronic periodontitis correlates with disease severity (7-11). We observed that EBV DNA was more frequently detected in patients with deeper periodontal pockets than in those with shallow ones or healthy controls (15, 16). We also observed that *P. gingivalis* was able to induce EBV reactivation through epigenetic regulation (17). Additionally, a large number of EBV-encoded small RNA (EBER)-positive B-cells were found in the gingival tissues of patients with chronic periodontitis (15). Thus, EBV is epidemiologically involved in the aetiology of chronic periodontitis. However, no causal relationship between EBV and chronic periodontitis has been delineated.

The level of gingival epithelial EBV infection is correlated with the severity of chronic periodontitis (18). EBV-infected cells reportedly express EBERs and EBV-encoded latent membrane protein (LMP1) (18). LMP1 is composed of 386 amino acids; it comprises a short *N*-terminal cytoplasmic domain, six transmembrane-spanning domains, and a *C*-terminal cytoplasmic tail (19-21), and mediates cell growth as an oncoprotein (19, 21). LMP1 expression is essential for EBV-mediated primary B-cell transformation in vitro and associated with a number of human malignancies (12, 13, 19, 21). In epithelial cells, LMP1 induces the expression of anti-apoptotic proteins and cell surface antigens (22-25). Additionally, it induces the production of matrix metalloproteinase 9, cyclo-oxygenase-2, and pro-inflammatory cytokines in immune cells (26-29). Taking these findings together, this viral protein activates inflammatory and immune-regulatory responses in EBV-infected cells.

Interleukin-8 (IL8), a potent neutrophil chemoattractant and activator, is a mediator of human chronic periodontitis *via* the accumulation and degranulation of neutrophils, which causes the subsequent destruction of periodontium (30, 31). Moreover, it strikingly upregulates the release of elastase by neutrophils, which significantly amplifies periodontal inflammation (32). IL8 thus plays a crucial role in the progression of chronic periodontitis.

In this study, to the best of our knowledge, we show for the first time that LMP1 attributable to its *C*-terminal activator regions (CTAR) domains in a human gingival epithelial cell line induces IL8 production *via* nuclear factor kappa B (NF- κ B) activation. Moreover, we discuss whether LMP1 in gingival epithelial cells possibly contributes to the progression of human chronic periodontitis.

Materials and Methods

Reagents and plasmids. Antibodies against phospho-I κ B α (Ser32) and phospho-p65 (Ser32) (Cell Signaling Technology, Danvers, MA, USA) and I κ B α , p65, β -actin, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) were purchased. A reporter plasmid expressing firefly luciferase under the control of NF- κ B (pGL3-5 $\times$ kB-luc) and dominant negative mutant of I κ B α (I κ B α Δ N) were used as

previously described (33). The *LMP1* expression vector (pSG-LMP1), its mutants, and control vector (pSG) (20) were generous gifts from Dr Martin Rowe (School of Cancer Sciences, University of Birmingham, UK).

Cell culture and transfection. The gingival epithelial cell line Ca9-22 was purchased from RIKEN BioResource Center (Tsukuba, Japan) and maintained at 37°C in Dulbecco's modified Eagle's medium (Sigma, St. Louis, MO, USA) with 10% heat-inactivated fetal bovine serum (Thermo Fisher Scientific, Rockford, IL, USA), penicillin, and streptomycin as previously described (34). Ca9-22 cells were transfected with pSG-LMP1 using Lipofectamine 2000 (Thermo Fisher Scientific), in accordance with the manufacturer's instructions.

mRNA preparation and real-time polymerase chain reactions (PCR). Experimental procedures for RNA purification and real-time (RT)-PCR were performed as previously described (34). For cDNA synthesis, total RNA was reverse transcribed using an RNA PCR kit (PrimeScript; Takara Bio, Shiga, Japan). The resulting cDNA mixture was subjected to RT-PCR analysis using SYBR Premix Ex Taq solution (Takara Bio) containing sense and antisense primers. The following primer sequences were used: *IL8*, forward (5'-CTTGTCATTGCCAGCTGTGT-3') and reverse (5'-TGACTGT GGAGTTTGGCTG-3'); *GAPDH*, forward (5'-TGCACCACC AACTGCTAGC-3') and reverse (5'-GGCATGGACTGTGGTCA TGAG-3'). PCR assays were performed using a TP-800 Thermal Cycler Dice Real-Time System (Takara Bio) and analysed using software provided by the device manufacturer. Thermal cycling conditions were 40 cycles at 95°C for 5 s, 60°C for 30 s, and 72°C for 1 min. All RT-PCR experiments were performed in triplicate; the specificity of each product was verified by melting curve analysis. The calculated level of gene expression was normalized to that of *GAPDH* mRNA.

IL8 measurements. IL8 in Ca9-22 cell-culture supernatants were measured using a human enzyme-linked immunosorbent assay (ELISA) kit for IL8 (R&D systems, Minneapolis, MN, USA), according to the manufacturer's instructions. All experiments were performed in triplicate, and data presented are representative of three independent experiments.

Preparation of cytoplasmic and nuclear extracts. Ca9-22 cells (2×10^5 cells/ml) were transfected with or without pSG-LMP1 for 24 h. Cells were washed with cold phosphate-buffered saline, resuspended in lysis buffer, and centrifuged. The supernatant was collected (whole-cell extract) and stored at -80°C until use. Precipitated cells were resuspended in cytoplasmic lysis buffer (Chemicon International, Temecula, CA, USA) and incubated for 15 min on ice. The cells were vortexed and centrifuged (10 min, $20,000 \times g$), and the supernatant (cytoplasmic extract) was removed. Pelleted cells were washed twice with cytoplasmic buffer to remove any trace of proteins, resuspended in nuclear lysis buffer (Chemicon International), and incubated for 15 min on ice. Lysed nuclei were sonicated for 10 s and centrifuged (15 min, $20,000 \times g$); the supernatant (nuclear extract) was stored at -80°C. The Pierce BCA Protein Assay-Reducing Agent Compatible kit (Thermo Fisher Scientific) was used for standardizing protein concentration.

Western blotting. Experimental procedures for western blotting were performed as previously described (34, 35). Briefly, equal amounts of protein (15 μ g) were separated by sodium dodecyl sulfate - poly

Figure 1. Latent membrane protein 1 (LMP1) promotes interleukin-8 (IL8) production in a human gingival epithelial cell line. Ca9-22 cells were transfected with pSG-LMP1 (0.2 μ g) or pSG (0.2 μ g) as control vector (ContV) for different times (A, C) and with pSG-LMP1 at different concentrations (0.05, 0.1, or 0.2 μ g) for 24 h (B, D). A, B: The cells were harvested and the level of IL8 mRNA was determined using reverse transcription polymerase chain reaction analysis with specific primers. C, D: IL8 released into the culture supernatants was determined using enzyme-linked immunosorbent assay. The values are presented as mean $\pm$ SD; n=3. **Significantly different at $p<0.0001$).

acrylamide gel electrophoresis and transferred to a polyvinylidene fluoride membrane (EMD Millipore Corporation, Billerica, MA, USA). The membrane was probed and visualised using a SuperSignal West Pico enhanced chemiluminescence kit (Thermo Fisher Scientific).

Transient luciferase assay. Ca9-22 cells (4×10^5 cells/ml) were transfected with 200 ng of reporter plasmids (pGL3-5 κ B-luc) and 10 ng of the internal control plasmid (pRL-TK) expressing Renilla luciferase (Promega, Madison, WI, USA), with or without LMP1 or I κ B α ΔN for 24 h, using Lipofectamine 2000. These cells were harvested using Passive Lysis Buffer (Promega); the extracts were assessed for luciferase activity using Dual-Luciferase Assay System (Promega) as previously described (17, 35).

Statistical analysis. Mean values $\pm$ standard deviation (SD) were calculated. Statistical analysis was performed using one-way analysis of variance with Tukey's multiple comparisons test; $p<0.05$ was considered to be statistically significant.

Results

LMP1 expression induces IL8 production in a human gingival epithelial cell line. RT-PCR analysis was used to investigate the effects of LMP1 on IL8 expression in Ca9-22 cells. LMP1 transfection induced significant IL8 expression. The IL8 mRNA level was up-regulated time-

Figure 2. Latent membrane protein 1 (*LMP1*) induces nuclear factor kappa B (*NF-κB*) activation. *Ca9-22* cells were transfected with *pSG-LMP1* or *pSG* (0.05 or 0.2 μg) as control vector (*ContV*). A: Proteins in the cytoplasmic extract (left panel) and the nuclear extract (right panel) were separated by polyacrylamide gel electrophoresis and then immunoblotted. Purity of the cytoplasmic and nuclear extracts was confirmed using antibodies specific for histone H3 as a representative of nuclear proteins and β-actin as a representative of cytoplasmic proteins. B: *Ca9-22* cells were transfected with *pSG-LMP1* (0.05, 0.1, or 0.2 μg) and 20 ng of the *pGL3-5κB-luc* reporter plasmid. The cells were harvested and luciferase (*Luc*) activity was measured 24 hours after transfection. The values are presented as mean±SD; n=3. Significantly different at **p*<0.05 and ***p*<0.0001.

independently (Figure 1A) and dose-dependently (Figure 1B) in response to *LMP1* transfection, but was not in cells transfected with the control vector. We next investigated the effects of *LMP1* on IL8 protein production. As shown in Figure 1C and D, extremely high concentrations of IL8 were produced due to *LMP1* in time- and dose-dependent manners. These results indicate that *LMP1* in human gingival cells may be a potent inducer of IL8 production.

***LMP1* activates *NF-κB* transcription in *Ca9-22* cells.** *NF-κB* is an inducible cellular transcription factor that regulates a variety of cellular genes involved in controlling inflammatory and immune responses (36). *NF-κB* normally binds to its inhibitor IκB present in the cytoplasm. Upon stimulation, intracellular signalling activates the IκB kinase complex, which sequentially phosphorylates two serine residues (Ser32/36) in IκBα (36). This results in the degradation of IκB by the 26S proteasome and consequent nuclear translocation of *NF-κB*.

As *NF-κB* activity is important in mediating IL8 production (36, 37), we next examined whether *LMP1* activated *NF-κB* in *Ca9-22* cells. We analysed the phosphorylation of *NF-κB* (p65) and IκBα using western blotting. As shown in Figure 2A, the level of phosphorylated IκBα increased, and that of IκBα conversely decreased due to *LMP1*. In parallel with this, phosphorylation increased at Ser536 of the *NF-κB* p65 subunit, which plays a key role in

the transcriptional competence of *NF-κB* and the nuclear translocation of p65 (36). To further examine whether *NF-κB* activation by *LMP1* in *Ca9-22* cells occurs at the transcriptional level, we performed luciferase assays using a reporter plasmid whose expression is proportional to *NF-κB* activity. As shown in Figure 2B, *LMP1* up-regulated *NF-κB* transcription in a dose-dependent manner. These results indicate that *LMP1* activates *NF-κB* in human gingival epithelial cells and that it may induce IL8 production.

***NF-κB* is involved in *LMP1*-induced IL8 production from *Ca9-22* cells.** We next examined whether *NF-κB* was involved in *LMP1*-induced IL8 production in *Ca9-22* cells. We used a dominant negative mutant of IκBα (IκBαΔN) to interfere with *NF-κB* activity (38). As shown in Figure 3A, *LMP1*-induced *NF-κB* activity was inhibited by IκBαΔN in a dose-dependent manner. In parallel with this, *LMP1*-induced IL8 production was also inhibited (Figure 3B). These results suggest that *NF-κB* is a potent regulator of *LMP1*-induced IL8 production from human gingival epithelial cells.

The CTAR1 and CTAR2 domains of *LMP1* contribute to *LMP1*-induced IL8 production from *Ca9-22* cells. Previous studies have identified two functional domains in the *LMP1* C-terminal cytoplasmic tail, CTAR1 (amino acids 187-231) and 2 (amino acids 351-386), which are important for *NF-*

Figure 3. The inhibitor of kappa B alpha ($I\kappa B\alpha$) mutant inhibits EBV-encoded latent membrane protein 1 (LMP1)-induced nuclear factor kappa B (NF- κ B) activation and interleukin-8 (IL8) production. Ca9-22 cells were transfected with $I\kappa B\alpha\Delta N$ -expressing plasmid (0.05 or 0.2 μ g), 0.2 μ g of pSG-LMP1, and 20 ng of pGL3-5 $\times$ κB-luc reporter plasmid for 24 h. A: Luciferase (Luc) assay was then performed. B: Ca9-22 cells were transfected with $I\kappa B\alpha\Delta N$ (0.05 or 0.2 μ g) and 0.2 μ g of pSG-LMP1 for 24 h. Enzyme-linked immunosorbent assay test for IL8 was then performed. The values are presented as mean $\pm$ SD; n=3. **Significantly different at $p<0.0001$.

κB activation (20, 25, 39). Proximal CTAR1 interacts with several tumor necrosis factor (TNF) receptor-associated factors (TRAFs) and induces NF-κB signalling (25, 39). Distal CTAR2 binds to the TNF receptor-associated death domain (TRADD) protein and mediates NF-κB signalling (20). Thus, we employed two *LMP1* mutants, *LMP1*Δ(187-351) lacking CTAR1 and *LMP1*Δ(349-386) lacking CTAR2, to analyse the effects of such mutations on NF-κB activation. The expression of *LMP1*Δ(187-351) or *LMP1*Δ(349-386) significantly reduced NF-κB activity (Figure 4A) and reduced IL8 production by approximately 80-90% compared with expression of wild-type *LMP1* (Figure 4B). Therefore, both TRAF-interacting CTAR1 and TRADD-interacting CTAR2 should be necessary for *LMP1*-induced IL8 production from human gingival epithelial cells.

Discussion

Although emerging evidence suggests an association between EBV and human chronic periodontitis, whose major aetiology has recently been revealed to be neutrophil infiltration into the periodontium, how EBV relates to the progression of periodontitis is not understood. To clarify this issue, it would be important to identify and characterize any complex pro-inflammatory cytokine network involved in periodontal pathogenesis and its link to EBV. We, therefore, first examined whether EBV *LMP1* in a human gingival cell line, Ca9-22, induced IL8 production. As expected, *LMP1* induced IL8 production. In addition, for IL8, the activation

Figure 4. C-terminal activator region 1 (CTAR1) and CTAR2 domains of latent membrane protein 1 (LMP1) are involved in LMP1-induced nuclear factor kappa B (NF-κB) activation and interleukin-8 (IL8) production. Ca9-22 cells were transfected with 0.2 μ g of the wild-type or the indicated mutant *LMP1* plasmids together with 20 ng of the pGL3-5 $\times$ κB-luc reporter plasmid for 24 h. A: Luciferase (Luc) assay was then performed as shown. B: Ca9-22 cells were transfected with wild-type or mutant (Δ) *LMP1* plasmids for 24 h. Cell culture supernatants were prepared and assayed for IL8 protein. The values are presented as mean $\pm$ SD; n=3. **Significantly different at $p<0.0001$.

of NF-κB was necessary. In parallel with this, mRNA expression of other NF-κB-driven pro-inflammatory cytokines, namely, IL1 β , IL6, and TNF α , was also up-regulated by LMP1 (data not shown).

Like other infectious diseases, adhesion to and subsequent invasion into the epithelia by pathogens are critical steps in the initiation of periodontitis. Bacterial adhesion to gingival epithelium reportedly induces the production of various immune-response mediators (1, 40). Among these, IL8, a potent neutrophil chemoattractant and activator, is such a mediator of human chronic periodontitis that acts *via* its involvement in the accumulation and degranulation of neutrophils, which causes subsequent destruction of the periodontium (30, 31). Many investigators have demonstrated the presence of IL8 in gingival crevicular fluid and pro-inflammatory cytokine levels in gingival crevicular fluid being closely associated with the severity of inflammation and periodontal destruction (41-44). Moreover, primed by IL1 β , IL8 strikingly up-regulates neutrophils for elastase release, giving rise to the significant increase of periodontal inflammation (32). IL8 is produced excessively or continuously in response to accumulated periodontopathic bacteria and their products in human gingival crevices or

periodontal pockets and is an important determinant of the progression of chronic periodontitis (2). In the present study, extremely high levels of IL8 were induced due to *LMP1* in Ca9-22 cells. This suggests that *LMP1* contributes to the progression of chronic periodontitis by promoting the production of pro-inflammatory cytokines, IL8 in particular.

Genes of several pro-inflammatory cytokines, including chemokines, contain NF- κ B-binding sites in their proximal promoters (36). Hence, NF- κ B activation possibly also plays a key role in virus-dependent cytokine expression and pathogenesis with inflammation. In the present study, *LMP1* stimulated simultaneous phosphorylation of I κ B α and NF- κ B p65 and the transcription of *NF- κ B*. Additionally, I κ B α Δ N inhibited *LMP1*-induced NF- κ B activation and subsequent IL8 production in a dose-dependent manner. This suggests that NF- κ B is a potent regulator of *LMP1*-induced IL8 production from human gingival epithelial cells. In this connection, this NF- κ B-dependent IL8 production has been reported to be associated with distinct human cells of the periodontium, such as a cervical squamous cell carcinoma cell line and umbilical vein endothelial cells (45, 46).

Regarding *LMP1*-related NF- κ B activation, the mutation analysis of *LMP1* has identified CTAR1 and CTAR2 as independent effectors of NF- κ B activation (20, 25, 39). CTAR1 binds to TRAF1, -2, -3, and -5 through a consensus TRAF-binding motif, while CTAR2 binds to the TRADD (20, 25, 39). In the present study, we found for the first time that *LMP1* Δ (187-351) and *LMP1* Δ (349-386), lacking CTAR1 and CTAR2, respectively, significantly reduced *LMP1*-induced IL8 production. Therefore, *LMP1*, particularly its CTAR domains, may be responsible for IL8 production from EBV-infected human gingival epithelial cells.

Taking our findings together, we confirmed that NF- κ B activation was responsible for *LMP1*-induced IL8 production by the gingival epithelial cell line. However, this was only obtained from a single cell line. Although Ca9-22 has been employed in many periodontal studies (47-49), it is important to confirm what we found in this study in other gingival epithelial cells to clarify the physiological relevance of our findings.

Chronic periodontitis and EBV infection in the periodontium quite often coincide, where high numbers of EBER-positive B-cells and EBER-positive gingival epithelial cells are observed in the gingival tissues of patients with periodontal disease (15, 18). Moreover, the involvement of EBV in the aetiology of human chronic periodontitis is supported by several observations: Anti-herpesvirus drug treatment reportedly resulted in an undetectable level of EBV with simultaneous periodontal improvement (10), and increased concentrations of pro-inflammatory cytokines in sera were consistently observed in EBV-infected patients (21, 40). In addition, as described in our previous report, butyric acid produced by

periodontopathic bacteria may cause EBV reactivation in the periodontium of EBV-infected individuals (17). Furthermore, the present study showed that *LMP1* in the human gingival epithelial cell line Ca9-22 markedly induced IL8 production attributable to NF- κ B activation.

Considering the observations mentioned above and the involvement of infiltrated neutrophils in the aetiology of human chronic periodontitis, it is suggested that inducible IL8 production by EBV *LMP1* in the human gingival epithelium contributes to human chronic periodontitis. Therefore, EBV, as well as periodontopathic bacteria, should be considered as therapeutic targets when treating human chronic periodontitis. Further studies are needed to determine whether EBV is as an independent aetiological agent of human chronic periodontitis or whether it acts in combination with periodontopathic bacteria. The development of new treatments and superior preventative methods should be supported by a better understanding of the role of EBV in the aetiology of human chronic periodontitis.

Conflicts of Interest

The Authors declare that there are no conflicts of interest in regard to this study.

Authors' Contributions

N.W., K.N., and R.K. performed the experiments, analysed the data, contributed reagents/materials/analysis tools, prepared figures, and reviewed drafts of the article. A.K., O.T., and T.K. contributed reagents/materials/analysis tools, analyzed the data, and reviewed drafts of the paper. H.S., M.T., Y.O., and S.S. contributed to discussion, analyzed the data, and reviewed drafts of the article. K.I. conceived and designed the experiments, performed the experiments, analysed the data, authored or reviewed drafts of the article.

Acknowledgements

The Authors thank Dr Martin Rowe for the *LMP1* plasmids. This work was supported by JSPS KAKENHI (No.16K11526), Sato Fund, Dental Research Center, Nihon University School of Dentistry, and a Nihon University Multidisciplinary Research Grant for 2017.

References

- 1 Hajishengallis G: Periodontitis: From microbial immune subversion to systemic inflammation. *Nat Rev Immunol* 15(1): 30-44, 2015. PMID: 25534621. DOI: 10.1038/nri3785
- 2 Darveau RP: Periodontitis: A polymicrobial disruption of host homeostasis. *Nat Rev Microbiol* 8(7): 481-490, 2010. PMID: 20514045. DOI: 10.1038/nrmicro2337
- 3 Holt SC and Ebersole JL: *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia*: The "red complex", a prototype polybacterial pathogenic consortium in periodontitis. *Periodontol* 2000 38(1): 72-122, 2005. DOI: 10.1111/j.1600-0757.2005.00113.x

4. Abusleme L, Dupuy AK, Dutzan N, Silva N, Burleson JA, Strausbaugh LD, Gamonal J and Diaz PI: The subgingival microbiome in health and periodontitis and its relationship with community biomass and inflammation. *ISME J* 7(5): 1016-1025, 2013. PMID: 23303375. DOI: 10.1038/ismej.2012.174
5. Ledder RG, Gilbert P, Huws SA, Aarons L, Ashley MP, Hull PS and McBain AJ: Molecular analysis of the subgingival microbiota in health and disease. *J Appl Environ Microbiol* 73(2): 516-523, 2007. PMID: 17085691. DOI: 10.1128/AEM.01419-06
6. Riep B, Edesi-Neu B, Claessen F, Skarabis H, Ehmke B, Flemmig TF, Bernimoulin JP, Göbel UB and Moter A: Are putative periodontal pathogens reliable diagnostic markers? *J Clin Microbiol* 47(6): 1705-1711, 2009. PMID: 19386852. DOI: 10.1128/JCM.01387-08
7. Contreras A, Nowzari H and Slots J: Herpesviruses in periodontal pocket and gingival tissue specimens. *Oral Microbiol Immunol* 15(1): 15-18, 2000. PMID: 11155159. DOI: 10.1034/j.1399-302x.2000.150103.x
8. Lu H, Zhu C, Li F, Xu W, Tao D and Feng X: Putative periodontopathic bacteria and herpesviruses in pregnant women: a case-control study. *Sci Rep* 6: 27796, 2016. PMID: 27301874. DOI: 10.1038/srep2779
9. Slots J, Saygun I, Sabeti M and Kubar A: Epstein-Barr virus in oral diseases. *J Periodontol Res* 41(4): 235-244, 2006. PMID: 16827715. DOI: 10.1111/j.1600-0765.2006.00865.x
10. Saygun I, Kubar A, Özdemir A and Slots J: Periodontitis lesions are a source of salivary cytomegalovirus and Epstein-Barr virus. *J Periodontol Res* 40(2): 187-191, 2005. PMID: 15733155. DOI: 10.1111/j.1600-0765.2005.00790.x
11. Konstantinidis A, Sakellari D, Papa A and Antoniadis A: Real-time polymerase chain reaction quantification of Epstein-Barr virus in chronic periodontitis patients. *J Periodontol Res* 40(4): 294-298, 2005. PMID: 15966906. DOI: 10.1111/j.1600-0765.2005.00796.x
12. Schmidt CW and Misko IS: The ecology and pathology of Epstein-Barr virus. *Immunol Cell Biol* 73: 489-504, 1995. PMID: 8713470. DOI: 10.1038/icb.1995.79
13. Thorley-Lawson DA and Gross A: Persistence of the Epstein-Barr virus and the origins of associated lymphomas. *N Engl J Med* 350(13): 1328-1337, 2004. PMID: 15044644. DOI: 10.1056/NEJMra032015
14. Ikuta K, Satoh Y, Hoshikawa Y and Sairenji T: Detection of Epstein-Barr virus in salivas and throat washings in healthy adults and children. *Microbes Infect* 2(2): 115-120, 2000. PMID: 10742683. DOI: 10.1016/S1286-4579(00)00277-X
15. Kato A, Imai K, Ochiai K and Ogata Y: Higher prevalence of Epstein-Barr virus DNA in deeper periodontal pockets of chronic periodontitis in Japanese patients. *PLoS One* 8(8): e71990, 2013. PMID: 23991022. DOI: 10.1371/journal.pone.0071990
16. Kato A, Imai K, Ochiai K and Ogata Y: Prevalence and quantitative analysis of Epstein-Barr virus DNA and *Porphyromonas gingivalis* associated with Japanese chronic periodontitis patients. *Clin Oral Investig* 19(7): 1605-1610, 2015. PMID: 29233156. DOI: 10.1186/s12903-017-0438-6
17. Imai K, Inoue H, Tamura M, Cueno ME, Inoue H, Takeichi O, Kusama K, Saito I and Ochiai K: The periodontal pathogen *Porphyromonas gingivalis* induces the Epstein-Barr virus lytic switch transactivator ZEBRA by histone modification. *Biochimie* 94(3): 839-846, 2012. PMID: 22178321. DOI: 10.1016/j.biochi.2011.12.001
18. Vincent-Bugnas S, Vitale S, Mouline CC, Khaali W, Charbit Y, Mahler P, Prêcheur I, Hofman P, Maryanski JL and Doglio A: EBV infection is common in gingival epithelial cells of the periodontium and worsens during chronic periodontitis. *PLoS One* 8(2): e80336, 2013. PMID: 24367478. DOI: 10.1371/journal.pone.0080336
19. McFarland ED, Izumi KM and Mosialos G: Epstein-Barr virus transformation: involvement of latent membrane protein 1-mediated activation of NF- κ B. *Oncogene* 18(49): 6959-6964, 1999. PMID: 10602470. DOI: 10.1038/sj.onc.1203217
20. Floettmann JE and Rowe M: Epstein-Barr virus latent membrane protein-1 (LMP1) C-terminus activation region 2 (CTAR2) maps to the far C-terminus and requires oligomerisation for NF- κ B activation. *Oncogene* 15(15): 1851-1858, 1997. PMID: 9362452. DOI: 10.1038/sj.onc.1201359
21. Klein G: Epstein-Barr virus strategy in normal and neoplastic B-cells. *Cell* 77(6): 791-793, 1994. PMID: 8004668. DOI: 10.1016/0092-8674(94)90125-2
22. Henderson S, Rowe M, Gregory C, Croom-Carter D, Wang F, Longnecker R, Kieff E and Rickinson A: Induction of BCL-2 expression by Epstein-Barr virus latent membrane protein 1 protects infected B-cells from programmed cell death. *Cell* 65(7): 1107-1115, 1991. PMID: 1648447. DOI: 10.1016/0092-8674(91)90007-1
23. Laherty CD, Hu HM, Opipari AW, Wang F and Dixit VM: The Epstein-Barr virus LMP1 gene product induces A20 zinc finger protein expression by activating nuclear factor κ B. *J Biol Chem* 267(34): 24157-24160, 1997. PMID: 1332946.
24. Dawson CW, Rickinson AB and Young LS: Epstein-Barr virus latent membrane protein inhibits human epithelial cell differentiation. *Nature* 344(6268): 777-780, 1990. PMID: 2158628. DOI: 10.1038/344777a0
25. Huen DS, Henderson SA, Croom-Carter D and Rowe M: The Epstein-Barr virus latent membrane protein-1 (LMP1) mediates activation of NF- κ B and cell surface phenotype via two effector regions in its carboxy-terminal cytoplasmic domain. *Oncogene* 10(3): 549-560, 1995. PMID: 7845680.
26. Muroso S, Inoue H, Tanabe T, Joab I, Yoshizaki T, Furukawa M and Pagano JS: Induction of cyclooxygenase-2 by Epstein-Barr virus latent membrane protein 1 is involved in vascular endothelial growth factor production in nasopharyngeal carcinoma cells. *Proc Natl Acad Sci USA* 98: 6905-6910, 2001. PMID: 11381123. DOI: 10.1073/pnas.121016998
27. Yoshizaki T, Sato H, Furukawa M and Pagano JS: The expression of matrix metalloproteinase 9 is enhanced by Epstein-Barr virus latent membrane protein 1. *Proc Natl Acad Sci USA* 95(7): 3621-3626, 1998. PMID: 9520415. DOI: 10.1073/pnas.95.7.3621
28. Hatzivassiliou E and Mosialos G: Cellular signaling pathways engaged by the Epstein-Barr virus transforming protein LMP1. *Front Biosci* 7: d319-d329, 2002. PMID: 11779697.
29. Eliopoulos AG, Dawson CW, Mosialos G, Floettmann JE, Rowe M, Armitage RJ, Dawson J, Zapata JM, Kerr DJ, Wakelam MJ and Reed JC: CD40-induced growth inhibition in epithelial cells is mimicked by Epstein-Barr virus-encoded LMP1: involvement of TRAF3 as a common mediator. *Oncogene* 13(10): 2243-2254, 1996. PMID: 8950992.
30. Bickel M: The role of interleukin-8 in inflammation and mechanisms of regulation. *J Periodontol* 64: 456-460, 1993. PMID: 8315568.

- 31 Takashiba S, Takigawa M, Takahashi K, Myokai F, Nishimura F, Chihara T, Kurihara H, Nomura Y and Murayama Y: Interleukin-8 is a major neutrophil chemotactic factor derived from cultured human gingival fibroblasts stimulated with interleukin-1 β or tumor necrosis factor α . *Infect Immun* 60(12): 5253-5258, 1992. PMID: 1452358.
- 32 Brandolini L, Sergi R, Caselli G, Boraschi D, Locati M, Sozzani S and Bertini R: Interleukin-1 β primes interleukin-8-stimulated chemotaxis and elastase release in human neutrophils *via* its type I receptor. *Eur Cytokine Netw* 8(2): 173-178, 1997. PMID: 9262966.
- 33 Asamitsu K, Yamaguchi T, Nakata K, Hibi Y, Victoriano AF, Imai K, Onozaki K, Kitade Y and Okamoto T: Inhibition of human immunodeficiency virus type 1 replication by blocking I κ B kinase with noraristeromycin. *J Biochem* 144(5): 581-589, 2008. PMID: 9519889. DOI: 10.1089/aid.1998.14.293
- 34 Imai K, Kamio N, Cueno ME, Saito Y, Inoue H, Saito I and Ochiai K: Role of the histone H3 lysine 9 methyltransferase Suv39 h1 in maintaining Epstein-Barr virus latency in B95-8 cells. *FEBS J* 281(9): 2148-2158, 2014. PMID: 24588869. DOI: 10.1111/febs.12768
- 35 Imai K, Togami H and Okamoto T: Involvement of histone H3 lysine 9 (H3K9) methyltransferase G9a in the maintenance of HIV-1 latency and its reactivation by BIX01294. *J Biol Chem* 285(22): 16538-16545, 2010. PMID: 20335163. DOI: 10.1074/jbc.M110.103531
- 36 Hayden MS and Ghosh S: Signaling to NF- κ B. *Genes Develop* 18(18): 2195-2224, 2004. DOI: 10.1101/gad.1228704
- 37 Mukaida N: Interleukin-8: An expanding universe beyond neutrophil chemotaxis and activation. *Int J Hematol* 72(4): 391-398, 2000. PMID: 11197203.
- 38 O'Brien DP, Oltz EM and Van Ness BG: Coordinate transcription and V(D)J recombination of the κ immunoglobulin light-chain locus: NF- κ B-dependent and -independent pathways of activation. *Mol Cell Biol* 17(7): 3477-3487, 1997. DOI: 10.1128/MCB.17.7.3477
39. Kaye KM, Devergne O, Harada JN, Izumi KM, Yalamanchili R, Kieff E and Mosialos G: Tumor necrosis factor receptor associated factor 2 is a mediator of NF- κ B activation by latent infection membrane protein 1, the Epstein-Barr virus transforming protein. *Proc Natl Acad Sci USA* 93(20): 11085-11090, 1996. PMID: 8855313. DOI: 10.1073/pnas.93.20.11085
- 40 Slots J: Herpesviral-bacterial interactions in periodontal diseases. *Periodontol* 2000 52(1): 117-140, 2010. PMID: 20017799. DOI: 10.1111/j.1600-0757.2009.00308.x
- 41 Payne JB, Reinhardt RA, Masada MP, DuBois LM and Allison AC: Gingival crevicular fluid IL-8: Correlation with local IL-1 β levels and patient estrogen status. *J Periodontal Res* 28(6): 451-453, 1993. PMID: 8254463.
- 42 Fitzgerald JE and Kreutzer DL: Localization of interleukin-8 in human gingival tissues. *Oral Microbiol Immunol* 10(5): 297-303, 1995. PMID: 8596673.
- 43 Özmeriç, N, Bal B, Baloş K, Berker E and Bulut Ş: Levels of interleukin-1 β and interleukin-8 in gingival crevicular fluids in adult periodontitis. *J Periodontol* 66(10): 852-859, 1995. PMID: 8537867. DOI: 10.1902/jop.1995.66.10.852
- 44 Jin L, Söder B and Corbet EF: Interleukin-8 and granulocyte elastase in gingival crevicular fluid in relation to periodontopathogens in untreated adult periodontitis. *J Periodontol* 71(6): 929-939, 2000. PMID: 10914796. DOI: 10.1902/jop.2000. 71.6.929
- 45 Yoshizaki T, Horikawa T, Qing-chun R, Wakisaka N, Takeshita H, Sheen TS, Lee SY, Sato H and Furukawa M: Induction of interleukin-8 by Epstein-Barr virus latent membrane protein-1 and its correlation to angiogenesis in nasopharyngeal carcinoma. *Clin Cancer Res* 7: 1946-1951, 2001. PMID: 11448908.
- 46 Xiong A, Clarke-Katzenberg RH, Valenzuela G, Izumi KM and Millan MT: Epstein-Barr virus latent membrane protein 1 activates nuclear factor- κ B in human endothelial cells and inhibits apoptosis. *Transplantation* 78(1): 41-49, 2004. PMID: 15257037. DOI: 10.1097/01.tp.0000129805.02631.ef
- 47 Hieke C, Kriebel K, Engelmann R, Müller-Hilke B, Lang H and Kreikemeyer B: Human dental stem cells suppress PMN activity after infection with the periodontopathogens *Prevotella intermedia* and *Tannerella forsythia*. *Sci Rep* 6: 39096, 2016. DOI: 10.1038/srep39096
- 48 Suzuki K, Sakiyama Y, Usui M, Obama T, Kato R, Itabe H and Yamamoto M: Oxidized low-density lipoprotein increases interleukin-8 production in human gingival epithelial cell line Ca9-22. *J Periodontal Res* 45(4): 488-495, 2010. DOI: 10.1111/j.1600-0765.2009.01263.x
- 49 Takeuchi H, Setoguchi T, Machigashira M, Kanbara K and Izumi Y: Hydrogen sulfide inhibits cell proliferation and induces cell-cycle arrest *via* an elevated p21 Cip1 level in Ca9-22 cells. *J Periodontal Res* 43(1): 90-95, 2008. PMID: 20412422. DOI: 10.1111/j.1600-0765.2009.01263.x

Received July 17, 2019

Revised July 26, 2019

Accepted July 31, 2019