

Amelioration of Amyotrophic Lateral Sclerosis in SOD1^{G93A} Mice by M₂ Microglia from Transplanted Marrow

MICHAEL W. EPPERLY¹, RENEE FISHER¹, LORA RIGATTI², SIMON WATKINS³, XICHEN ZHANG¹, WEN HOU¹, DONNA SHIELDS¹, DARCY FRANICOLA¹, HULYA BAYIR⁴, HONG WANG¹, STEPHANIE THERMOZIER¹, ANDREW HENDERSON¹, CHRISTOPHER DONNELLY⁵, PETER WIPF⁶ and JOEL S. GREENBERGER¹

¹Department of Radiation Oncology, UPMC Hillman Cancer Center, Pittsburgh, PA, U.S.A.;

²Division of Laboratory Animal Resources (DLAR), University of Pittsburgh, Pittsburgh, PA, U.S.A.;

³Department of Cell Biology and Center for Images, University of Pittsburgh, Pittsburgh, PA, U.S.A.;

⁴Department of Critical Care Medicine, University of Pittsburgh, Pittsburgh, PA, U.S.A.;

⁵Department of Neurobiology, University of Pittsburgh, Pittsburgh, PA, U.S.A.;

⁶Department of Chemistry, University of Pittsburgh, Pittsburgh, PA, U.S.A.

Abstract. *Background/Aim:* The cause of fatal neuromuscular amyotrophic lateral sclerosis (ALS) is not known. *Materials and Methods:* Ninety-day-old superoxide-dismutase-1^{G93A} (SOD1^{G93A}) mice demonstrating level 1 paralysis, received 9.0 Gy total body irradiation (TBI) from a cesium source at 340 cGy per minute, and intravenous transplantation with 1×10⁶ C57BL/6 green fluorescent protein (GFP)+ donor bone marrow cells. *Results:* Paralysis-free survival was prolonged in TBI and bone marrow-transplanted SOD1^{G93A} mice from 100 to over 250 days (*p*=0.0018). Other mice transplanted with SOD1^{G93A} marrow or marrow treated with the free-radical scavenger MMS350 showed no therapeutic effect. GFP+ macrophage-2 (M₂) microglial cells of bone marrow origin, were seen at sites of degenerating anterior horn motor neurons. SOD1^{G93A} mice had a disruption in the blood-brain barrier permeability which was reversed by marrow transplant from C57BL/6 mice. SOD1^{G93A} marrow showed unexpected robust hematopoiesis in vitro, and radioresistance. *Conclusion:* After TBI, M₂ microglial cells from transplanted donor marrow extended the paralysis-free interval in SOD1^{G93A} mice.

Amyotrophic lateral sclerosis (ALS) is a fatal neuromuscular disease caused by progressive neurodegeneration of the motor neurons in the brain and spinal cord (1, 2). While the cause of ALS is unknown, 10-15% of patients have a familial history of disease (3-7), while 85-90% develop the disease sporadically. There are nearly 40 genetic mutations associated with ALS, the second most common of which is a mutation in the superoxide dismutase-1 (SOD1) gene (3-30). The SOD1^{G93A} mouse model phenotypically recapitulates ALS in human patients and exhibits progressive muscle wasting, paralysis, and death (18-25). Existing therapeutic interventions only modestly extend the lifespan of patients with ALS. Recently, approved antioxidant therapies are believed to slow disease progression. Experimental therapeutic strategies, including administration of antioxidant drugs, biological response modifiers, cytokine inhibitors, cytokine receptor antagonists, and introduction into the spinal cord of neurotrophic growth factors or mesenchymal stem cells engineered to produce neuroleukines (31-46), have met with incomplete success (2, 32). Marrow transplant strategies, which were designed to target specific defects (47-56), have been either ineffective or of limited success in ALS (51, 52).

The histopathology of ALS-affected spinal cord demonstrates gliosis, including the accumulation of both inflammatory macrophage-1 (M1) and neurotrophic factor-producing (M₂) microglia at the site of neurodegeneration (8). However, the pathophysiology of homing of these cells to the site of motor neurons, and their mechanism of action is not known (8-13, 15, 19-22). Furthermore, the origin of microglial cells in the thoracolumbar spinal cord, where ALS typically begins the process of ascending muscle functional loss is not known. A summary of recent research suggests that microglial cell accumulation at the site of degenerating

This article is freely accessible online.

Correspondence to: Joel S. Greenberger, MD, Professor and Chairman, Department of Radiation Oncology, UPMC Hillman Cancer Center, UPMC Cancer Pavilion, Rm. 533, 5150 Centre Avenue, Pittsburgh, PA 15232, U.S.A. Tel: +1 4126473602, Fax: +1 4126476029, e-mail: greenbergerjs@upmc.edu

Key Words: Total body irradiation, ALS, marrow origin, spinal cord M2 cells.

neurons represents multiple phases in the degenerative process, one inflammatory potentially initiating the disease process (28), and a second response to degenerating neurons with a production of repair-related cytokines (9, 10). There remains controversy over the interaction of glial cells with motor neurons in the development of paralysis (57, 58).

In the present studies, we tested the hypothesis whether cells of donor bone marrow origin might provide a therapeutic effect if delivered to SOD1^{G93A} mice after total body irradiation (TBI).

Materials and Methods

Mice and animal care. SOD1^{G93A} and control SOD1 transgenic mice containing 4 copies of normal human SOD1 gene were obtained from the Jackson Laboratories, Bar Harbor, Maine. SOD1^{G93A}-green fluorescent protein (GFP⁺) mice were bred according to published methods used to breed SMAD family member 3 (*Smad3*)^{-/-} Fanconi anemia group D2 (*Fancd2*)^{-/-} mice (59). Mice were derived from breeding pairs of SOD1^{G93A} and control mice. Female mice were housed at five mice per cage and male mice four per cage according to institutional regulations. Mice were fed standard Purina laboratory chow. All work was performed under protocol 18022000, approved by the University of Pittsburgh Institutional Care and Use Committee. Veterinary care was provided by the Division of Laboratory Animal Resources at the University of Pittsburgh.

Paralysis was scored as described by Hatzipetros *et al.* (26). Mice were euthanized when reaching paralysis with neurological score (NS) of 3. At NS 3, the mice still have complete use of their front legs but show paralysis in the rear leg. Mice with NS 1 have no paralysis but exhibit some trembling in the rear legs and a collapsing of the rear legs toward the lateral mid line when picked up. NS 2 paralysis reflects the beginning of paralysis in the rear legs with a complete collapse of the rear legs to the lateral midline when picked up; mice begin to show an altered gait but are still able to move easily around in the cage.

Mice were given no irradiation, 9 Gy, or 7.0 Gy TBI using a Shephard Mark I ¹³⁷Cs γ -ray source (J. L. Shepherd, San Fernando, CA, USA), according to published methods (59). Other mice received 9 Gy cranial spinal irradiation. Subgroups were given 10⁶ cells from B6 GFP⁺ or wild-type B6 mouse marrow intravenously after fractionated TBI, as described elsewhere (60).

TBI and craniospinal irradiation. Female and male mice received 9.0 Gy TBI from a cesium source at 340 cGy/min using a Shephard Mark I irradiator (J.L. Shepherd, San Fernando, Ca, USA). Bone marrow was isolated from C57BL/6 GFP⁺ mice (Jackson Laboratories, Bar Harbor, ME, USA). The irradiated SOD1^{G93A} mice were injected intravenously with 1×10⁶ C57BL/6 GFP⁺ donor bone marrow cells. Chimerism was determined from the presence of GFP⁺ cells in peripheral blood according to (60). The mice were followed for the development of NS 3 paralysis at which time they were sacrificed and the spinal cord was isolated and fixed in 2% paraformaldehyde (PFA). Some of the spinal cords were analyzed by single photon, confocal, ribbon-scanning microscopy for quantitation of bone marrow origin cells. Other spinal cords were fixed in 2% PFA, sectioned, and stained for M₁ and M₂ microglial cells.

Control C57BL/6NTac mice were transplanted at the same age with SOD1^{G93A} ALS bone marrow under the same conditions and followed for development of paralysis. To determine if spinal cord irradiation alone without bone marrow transplantation altered the development of paralysis, SOD1^{G93A} mice were irradiated with 9.0 Gy to the spinal cord and brain while shielding the remainder of the body. All mice were followed for the development of paralysis. In a separate group of SOD1^{G93A} mice, marrow transplant using B6 GFP⁺ or SOD1^{G93A} GFP⁺ 1×10⁶ bone marrow cells was performed at 24 h after TBI.

To determine if TBI alone delay paralysis, SOD1^{G93A} mice were irradiated to 7.0 Gy TBI without bone marrow transplantation. Mice were followed for development of paralysis.

Treatment with MMS350. The water-soluble radiation mitigator MMS350 was produced in the laboratory of Dr. Peter Wpif at the University of Pittsburgh, Pittsburgh, PA, USA and was administered at 400 mg/ml to mice in drinking water as described (61, 62) over days 60 until death from paralysis.

Continuous bone marrow culture. Long-term bone marrow cultures (LTBMCs) were established from the femur and tibia marrow of SOD1^{G93A} or control transgenic mice, as described previously (63). The contents of a femur and tibia (N=6/genotype) were flushed into McCoy's 5A medium (Gibco, Gaithersburg, MD, USA) supplemented with 25% horse serum (Cambrex, Rockland, ME, USA) and 10⁻⁵ M hydrocortisone sodium hemisuccinate. Cultures were incubated at 33°C in 7% CO₂. After 4 weeks, horse serum was replaced with 25% fetal bovine serum (FBS) (Gibco, Gaithersburg, MD, USA) (63). The cultures were observed weekly for hematopoietic cell production and cobblestone island formation. The number of cobblestone islands of 50 cells or more were scored weekly in each flask (63-64). A two-sided two-sample *t*-test was used to compare the number of cobblestone islands between culture groups each week. *p*-Values less than 0.05 were regarded as significant.

The cultures were observed weekly for hematopoietic cell production and adherent cell layer confluence. Non-adherent cell production data were expressed as the mean±standard error standard error of the mean (SEM) of weekly non-adherent cell number and cumulative non-adherent cell production. Confluence data were expressed as the mean±SEM of the percentage adherent cell layer confluence.

Establishment of clonal bone marrow stromal cell lines. Adherent cell layers from one 4-week-old LTBMCM from SOD1^{G93A} (63-64) and control mice were trypsinized and expanded by passage into Dulbecco's Modified Eagle's Medium (DMEM) with 10% FBS to establish bone marrow stromal cell lines according to published methods (63). Cells were passaged for 10 weeks to establish cell lines. Culture were incubated at 37°C in 5% CO₂.

Hematopoietic colony-forming cell lines and fresh marrow colony assays. Each week after explant and establishment of LTBMCMs, 1×10⁵ nonadherent cells were plated in triplicate in methylcellulose-containing medium, as previously published (63). Cells were incubated at 37°C in 5% CO₂. At day 7 after plating, colonies of ≥50 cells were counted. The plates scored at day 7 were returned to the incubator for scoring at day 14. Data for days 7 and 14 are presented as the mean±SEM of weekly colony-forming cells and cumulative colony-forming cells (63).

Table I. Source of antibodies and target cell identifiers.

Primary antibody target	Known functions of target
Glial fibrillary acidic protein (Novus Biologics, NB300-141)	Intermediate filament expressed by numerous cell types in the central nervous system including astrocytes and ependymal cells.
Allograft inflammatory factor 1 (Abcam, ab5076)	Participates in membrane ruffling and phagocytosis in activated microglia.
Neurofilament heavy polypeptide (Abcam, ab8135)	Provides structural support to axons and regulate axon diameter.
Neurotracer (ThermoFisher, N21482)	A fluorescent tracer that adheres to neurons
Oligo2 (Novus Biologics, NBP1-28667)	Oligodendrocyte transcription factor found in motor neurons and oligodendrocytes
Platelet endothelial cell adhesion molecule (Abcam, ab134168)	Involved in leukocyte transmigration, angiogenesis and integrin activation.

Establishment of bone marrow stromal cell lines. Adherent cell layers from one LTBM per treatment group were trypsinized and expanded by passage into DMEM supplemented with 10% FBS to establish bone marrow stromal cell lines according to published methods (63). Established criteria were used for authentication of cell lines (63). Genotyping of all bone marrow stromal cell lines established they were indeed from the SOD1^{G93A} or control genotype. The established lines maintained stromal cell features including capacity for osteoblast differentiation, and support of hematopoietic cells in co-culture. We documented cell surface phenotype, biology, and absence of capacity to differentiate into hematopoietic cells. These cell lines were characterized as bone marrow stromal cells or mesenchymal stem cells (63). Cell lines were incubated at 37°C in 5% CO₂ and passaged for 10 weeks to establish cell lines.

Establishment of single-cell-derived clonal marrow stromal cell lines. Single-cell cloning experiments were performed using fresh marrow, an adherent cell layer from a LTBM at week 13 after explant, and bone marrow stromal cell lines that were established as described (63). Flow cytometry was used to sort single cells into the wells of a 96-well plate. Cells were grown in DMEM supplemented with 20% FBS, 1% antibiotic-anti-mitotic solution, and 1% L-glutamine. Plates were observed weekly for growth. Clonal lines were established from wells that showed growth after single-cell plating.

Histochemistry and ribbon scanning, confocal single-photon microscopy. The detailed methods for the microscopy techniques of the spinal cord have been published in the web-based textbook (65).

Microbead assay for blood-brain barrier permeability. Blood-brain barrier permeability was analyzed by intravenous perfusion with 0.2 µm Fluorospheres (F8810; Invitrogen) on day 125 in control SOD1^{G93A} mice, SOD1^{G93A} mice transplanted on day 90 with GFP+ C57BL6 bone marrow and C57BL/6 mice.

Immunofluorescence assay for detection of astrocytes, neurons, glial cells, and immunocytes. Spinal cord was removed from control C57BL/6 mice, and SOD1^{G93A} mice with phase 1, phase 2 and phase 3 paralysis. The spinal cords were fixed in 2% paraformaldehyde (PFA; Thermo Fisher Scientific, Waltham, MA, USA), sectioned, and stained with antibodies to astrocytes, neurons, glial cells and immunocytes. The antibodies used, their vendor/supplier and cellular target function are shown in Table I. Secondary antibodies used include goat anti-rabbit and donkey anti-goat (Thermo Fisher Scientific, Waltham, MA, USA). The antibodies specific for astrocytes,

motor neurons, M₁ and M₂ microglial cells, endothelial cells, and hematopoietic cells have been described elsewhere (65). Cells were incubated with primary antibody mixture overnight at 4°C then washed with PBS three times for 5 min each. Secondary antibody solution was prepared at manufacturer-suggested dilutions in PBS and added to cells for 1 hour at room temperature. After washing with PBS three times for 5 min each, 0.5 µg/ml 4',6-diamine-2'-phenylindole dihydrochloride (DAPI; Millipore Sigma, St. Louis, MO, USA) in PBS was added for 10-20 min at room temperature to label nuclei. Cells were mounted and allowed to dry for 30 min. Pictures were taken using fluorescence microscopes at the University of Pittsburgh Center for Biologic Imaging. For each sample, three fields were subjected to analysis. Data are presented as the mean±standard error of the percentage of cells staining positively for the target protein.

Statistical analysis. For *in vivo* experiments, the mean and standard error of the mean for each group was determined and graphed using GraphPad Prism (GraphPad Software, LaJolla, CA, USA) to compare experimental groups. Student's unpaired *t*-test was used to compare data of each group at different time points to those for the control group, as well as to compare all days for each experimental group (including the control) to day 0 (non-irradiated baseline) of the control group.

For LTBMcs, GraphPad Prism (GraphPad Software) was used for statistical analysis and plotting graphs. Graphs were plotted using the mean±SEM. The data were analyzed using two-tailed Student's *t*-test. In this exploratory experiment, *p*-values were not corrected for multiple comparisons. Differences were considered significant at *p*<0.05 (27). At each week of LTBM per each of the endpoints (*i.e.* weekly non-adherent cells per flask, percent confluence, day-7 colonies, and day-14 colonies), data were summarized with mean±standard deviation. Comparisons were made using the one-way analysis of variation (ANOVA) F-test at each time point, followed by Tukey's multiple comparisons.

The description of calculation of D₀, which is the irradiation dose required to reduce survival to 37% on the linear portion of the survival curve, and \bar{n} , which represents the shoulder on the survival curve as calculated by the back extrapolation of the linear portion of the survival curve to the y axis, for radiation survival curves has been published in detail elsewhere (63).

Results

Establishment of bone marrow chimeric GFP+ SOD1^{G93A} mice. After marrow transplant, as described in the Materials




Figure 1. Effect of total body irradiation (TBI) and bone marrow transplantation (BMT) on paralysis-free interval. Superoxide dismutase-1 (SOD1)^{G93A} mice at 90 days after birth received total body irradiation to 9 Gy. A: Mice were injected intravenously with 1×10^6 freshly removed tibia and femur bone marrow from C57BL/6-green fluorescent protein (GFP)+ mice and followed for development of stage 3 paralysis at which time the mice were sacrificed. Non-treated SOD1^{G93A} mice had an average survival of 129 ± 3 days ($n=5$). In contrast, TBI-mice which received bone marrow transplant had a significant increase in mean survival to 233 ± 23 days ($p=0.0045$) ($n=6$). Control wild-type C57BL/6 mice ($n=10$) were irradiated to 10 Gy and transplanted with bone marrow from SOD1^{G93A} mice and followed for development of paralysis. No paralysis, as suggested in several recent publications (21-24) was detected up to 280 days following transplantation ($p=0.0018$). B: Controls included mice which received 9.0 Gy brain and spine irradiation ($n=5$), sub-lethal TBI of 7.0 Gy alone ($n=5$), radiation mitigator MMS350 in drinking water ($n=4$), or transplant of SOD1^{G93A} marrow ($n=4$). The horizontal line for each group represents the average age at the time of sacrifice.

and Methods, blood samples were checked for chimerism at day 60 and 90. Mice with over 80% donor origin GFP+ cells were considered to be successfully transplanted.

Quantitation of TBI and bone marrow transplantation delayed hind limb paralysis. TBI and marrow transplant significantly delayed paralysis and extended survival of SOD1^{G93A} mice (Figure 1A). TBI plus normal marrow transplant but not sub-TBI, reduced TBI dose, transplant of SOD1^{G93A} donor marrow nor administration of MMS350 prolonged the paralysis-free interval ($p < 0.0039$) (Figure 1B).

The stages of paralysis are shown histologically in Figure 2. Bone marrow-derived M₂ microglia were found in the spinal cords of SOD1^{G93A} mice transplanted with GFP+ donor marrow at 120 days of age when other groups are developing stage 3 paralysis (Figure 3). GFP+ cells were seen throughout the spinal cord (Figure 4). The blood-brain barrier was permeable to microbeads in non-irradiated and non-transplanted SOD1^{G93A} mice (Figure 5) confirming findings of a prior publication (47). In contrast, TBI-treated, then GFP+ marrow-transplanted mice showed correction of the blood-brain barrier defect (Figure 5). Control wild-type mice transplanted with SOD1^{G93A} bone marrow demonstrated no paralysis (Figure 1A). There was no transfer of cells of SOD1^{G93A} GFP+ donor bone marrow origin to the spinal cord of wild-type recipients. The blood-brain barrier was no longer disrupted in TBI-treated normal marrow-transplanted SOD1^{G93A} mice (Figure 5).

The results show that TBI and bone marrow transplantation significantly extended the survival of SOD1^{G93A} mice, and was a robust neuroprotective and translatable strategy to prevent motor neuron degeneration starting at the time of NS 1 paralysis in SOD1^{G93A} mice (Figure 1).

Analysis of subsets of cells of bone marrow origin in spinal cord of TBI-treated and bone marrow-transplanted SOD1^{G93A} mice. We identified microglia from GFP+ marrow origin in the spinal cords of SOD1^{G93A} TBI/marrow-transplanted mice (Figures 3 and 4). We tested whether there were GFP+ M₂ microglial cells in the anterior horns of the spinal cords at days 120, 200, and 280 in GFP+ marrow-transplanted SOD1^{G93A} mice (Figure 3). The spinal cord showed significant infiltration with GFP+ cells of marrow origin at multiple levels (Figures 3 and 4). We did not see cells of marrow origin in the spinal cords of TBI-treated wild-type C57BL/6 mice that were transplanted with GFP+ SOD1^{G93A} marrow.

Demonstration of reversal of blood/brain barrier permeability in SOD1^{G93A} mice. SOD1^{G93A} mice demonstrated the blood-brain barrier was permeable to 2- μ m diameter beads (Figure 5), confirming previous findings (47). In contrast, control C57BL/6 mice and C57BL/6 mice that




Figure 2. Histopathological stages of spinal cord motor neuron loss in superoxide dismutase-1 ($SOD1^{G93A}$) mice (H&E $\times 20$). With the advancement of each stage, there was increased degeneration and loss of motor neurons, astrogliosis in the areas of motor neuron loss, and vacuolation within the gray matter ($n=3$ at each stage). Lesions were more pronounced in the ventral horns (bar= $100\ \mu\text{m}$).




Figure 3. Histochemical detection of M_2 microglial cells of bone marrow origin in the spinal cord of superoxide dismutase-1 ($SOD1^{G93A}$) mice. A: Cross section of spinal cord at recipient age 120 days which is 30 days post irradiation and transplantation of green fluorescent protein (GFP)+ bone marrow cells ($n=3$). Labelling is glial fibrillary acidic protein (GFAP) (red), nuclei (blue), ionized calcium-binding adapter molecule 1 (white), and stem cells (green). There is almost complete co-localization between the green cells of marrow origin and the white M_2 microglia ($\times 100$). B and C: Boxed inset region from A showing M_2 microglia (B) overlaying exactly with the GFP-labeled cells (C), highlighted with yellow arrows. Bar= $500\ \mu\text{m}$ ($\times 500$).

had been transplanted with $SOD1^{G93A}$ marrow showed no such detectable blood–brain barrier permeability (Figure 5).

Failure of isolated craniospinal irradiation and of sub-lethal TBI with no marrow transplant to prolong the paralysis-free interval in $SOD1^{G93A}$ mice. As shown in Figure 1B,

$SOD1^{G93A}$ mice treated with 9.25 Gy to the brain and spinal cord only while shielding the rest of the body, and those receiving 7.0 Gy TBI without marrow transplant showed no therapeutic effect. Furthermore, $SOD1^{G93A}$ mice treated with radiation mitigator MMS350 in drinking water starting at day 60 showed no therapeutic effect (Figure 1B).




Figure 4. Rendered three-dimensional image of superoxide dismutase-1 ($SOD1^{G93A}$) murine spinal cord following irradiation and transplantation of green fluorescent protein (GFP)+ bone marrow cells. Images were collected following CUBIC clearing and imaging using a Caliber XT ribbon scanning confocal microscope (25×1.15 NA water objective, WD 2.4 mm) ($n=3$). A: En face reconstruction of the raw image. GFP+ cells can be seen throughout the tissue, but are more concentrated in the lumbar region in the lateral nerve branches. B: The GFP+ cells in the same image set in which the positive structures are highlighted as green spheres. C: The 3D nature of the reconstruction is clearly visible. Bar=1 cm. (Link to movie of Figure 4).




Figure 5. The blood–brain barrier is permeable in superoxide dismutase-1 ($SOD1^{G93A}$) mice. A: Section through spinal cord of an ALS mouse (age 100 days) perfused with $0.2\ \mu\text{m}$ Fluospheres (F8810, Invitrogen) shown in red ($n=3$). The actin cytoskeleton is green and blue defines the nuclei (bar= $250\ \mu\text{m}$). Beads are present throughout the soma of the spinal cord (Panel B). However, in TBI/transplanted mouse at 30 days post transplantation (Panel C) no beads are present in the spinal cord.

Hematopoiesis in LT BMCs from $SOD1^{G93A}$ mice. We tested the effect of the $SOD1^{G93A}$ genotype on limiting the duration of hematopoiesis in LT BMCs as a marker of defective antioxidant capacity according to prior publications

(59, 61-64). Unexpectedly, $SOD1^{G93A}$ mouse marrow had greater longevity than did that from $SOD1$ transgenic control mice. Stability of the adherent layer was equivalent in $SOD1^{G93A}$ and $SOD1$ transgenic control mice (Figure 6A).

Table II. Radioresistance of superoxide dismutase-1 (SOD1)^{G93A} isolated bone marrow and bone marrow stromal cells.

Mouse Strain	SOD1 ^{G93A} bone marrow cells		SOD1 ^{G93A} bone marrow stromal cells	
	Do (Gy)	n	Do (Gy)	n
SOD1 Transgenic	7.583±0.819	1.554±0.05989	8.564±1.564	1.319±0.0799
SOD1 ^{G93A}	4.791±0.5162	1.593±0.01132	3.803±0.3985	1.653±0.07226
p-Value	0.0488		0.0210	0.0228

Stem cell islands, cobblestones, were significantly greater in number (Figure 6B). There was a significantly greater production of total hematopoietic cells (Figure 6C), and greater duration of production of total cell forming day 7 (Figure 6D) and day 14 (Figure 6E) colony-forming progenitors (Supplemental Tables I-V may be found at <https://pitt.box.com/s/ajzj6ov2cw9ttrbnsfnfz3p7d7so7mob>).

Radiation resistance of clonal bone marrow stromal cell lines from SOD1^{G93A} mice. In prior studies, greater longevity of hematopoiesis in LTBMcs correlated with radioresistance of hematopoietic progenitors, suggesting a greater capacity of cells to tolerate oxidative stress (63). The data revealed that both SOD1^{G93A} mouse isolated bone marrow CFU-GEM (Figure 7A, Table II) and marrow culture-derived stromal cell lines (Figure 7B, Table II) were radioresistant.

Number of motor neurons in thoracic and lumbar spinal cord of SOD1^{G93A} mice is reduced compared to controls. The increased number of hematopoietic progenitor cells in LTBMcs and radioresistance of both fresh marrow colony-forming units-granulocyte, erythrocyte, monocyte, megakaryocyte and stromal cell lines from SOD1^{G93A} mice suggested that the defective *SOD1* gene may not have reduced stem cell numbers in the marrow, but may have depleted the critical motor neurons in the spinal cord.

We counted motor neurons in spinal cords from mice of the SOD1^{G93A} genotype in NS 0, 1, 2, 3. While the number of motor neurons was reduced at NS 1, 2, and 3 compared to stage 0, there was no clear direct stage-specific loss of motor neurons with progression of paralysis (Figure 8).

Discussion

In the present study with SOD1^{G93A} ALS mice, we demonstrated that TBI and bone marrow transplantation during the symptomatic stage of the disease (day 90) exerted a profound therapeutic effect by delaying paralysis, and extending survival two-fold from 120 to over 250 days. This treatment regimen was significantly more effective than previous mesenchymal stem cell or bone marrow

transplantation studies in SOD1^{G93A} mice, and in other models of ALS (51-52). The present treatment paradigm more accurately recapitulates a therapeutic approach in patients with ALS since they often exhibit symptoms at the time of diagnosis. In other studies, from our laboratory using marrow transplant at day 60 before symptoms, we observed that the prolongation of survival was even better (data not shown). We found that SOD1^{G93A} mice, but not transgenic mice with four copies of normal human *SOD1* displayed permeability of the blood-brain barrier, which was reversible by TBI and marrow transplant, and we identified bone marrow-derived M₂ microglia at the site of spinal motor neurons.

The present data suggest that the observed therapeutic effect of TBI and marrow transplant is mediated by M₂ microglia of marrow origin. In the present experiments, TBI and marrow transplant producing chimeric SOD1^{G93A} mice resulted in M₂ microglial cells of bone marrow origin accumulating at the site of anterior horn cell degeneration. Mice receiving 9.0 Gy craniospinal irradiation or sub-lethal 7.0 Gy TBI irradiation alone did not show a therapeutic effect. Continuous antioxidant treatment with MMS350 (61,62) in drinking water also showed no beneficial effect.

Our data are the first to be reported of successful TBI and bone marrow transplantation in symptomatic SOD1^{G93A} mice. Treatment did delay paralysis and more than doubled the lifespan. We determined that sub-lethal TBI, cranial spinal, or of the spinal cord only was not therapeutic. A marrow cell population facilitated accumulation of therapeutic M₂ microglial cells of marrow origin in the spinal cord at the site of motor neuron, but only after TBI. These studies are the first to demonstrate that ionizing irradiation in TBI is therapeutic for ALS.

Our novel SOD1^{G93A} GFP+ mice and our unique single-photon, confocal ribbon-scanning microscopy allowed us to count all green motor neurons in the entire spinal cord. We have constructed a jig which allows non-anesthetized immobilized mice to be irradiated in specific areas of the spinal cord with precise radiation doses. In this way, less than 1% of the prescribed dose is delivered outside the irradiated field using our Stereotactic Radiosurgery Unit at our TrueBeam/Varian linear accelerator. The combination of




Figure 6. Continued

these novel techniques allowed us to quantitate the effect of each of the treatment modalities on preserving the number of intact and non-degenerating anterior horn motor neuron cells. Only TBI plus bone marrow transplantation facilitated the therapeutic effect in these SOD1^{G93A} mice.

TBI for preparation of recipients for marrow transplant has been used for over 60 years in the treatment of not only

cancer, but also autoimmune diseases and genetic disorders, including some that affect the central nervous system (49, 50). Use of bone marrow transplant in ALS has not been successful since results showed either no effect (51), or minimal effect (52). Irradiation effects on the spinal cord and the radiobiology of neurons have not been considered as a possible therapeutic modality for diseases of motor neurons in ALS (53). The role




Figure 6. Improved hematopoiesis in long-term bone marrow cultures from superoxide dismutase-1 (*SOD1*^{G93A}) mice compared to control *SOD1* transgenic mice containing four copies of the normal human transgene (*SOD1* transgene) ($n=4$). Absolute and cumulative data for surface area confluence (A), cobblestone islands per flask per week (B), non-adherent cells produced per flask per week (C), and colony-forming cells generated per flask per week at day 7 (D) and day 14 (E). All statistical analysis for the data is shown in Supplementary Tables I to V (<https://pitt.box.com/s/ajzj6ov2cw9ttrbnsfnfz3p7d7so7mob>).

of cells of marrow origin, both pro-inflammatory and anti-inflammatory, in the clinical evolution of ALS has not been mechanistically coupled with the role of irradiation relative to abnormalities in the blood–brain barrier in ALS, nor have the effects of transplanted marrow cells into patients with ALS been studied relative to prolongation of the paralysis-free interval (36, 41, 47, 54-58, 66-69).

Recent data with *SOD1*^{G93A} mice suggest that the inflammasome and significantly, interleukin 1 family cytokines, may mediate motor neuron cell death *via* induction of radical oxygen species pathways (70-78). How M₂ microglial from marrow might neutralize these interactions at the level of the microglial cell to motor neuron interface is unknown. Because 10% of ALS cases are familial, these *SOD1*^{G93A} mice may represent the most appropriate genetic model for future ALS studies.

The mechanism of cell death in ALS motor neurons is not known. Apoptosis, necroptosis, ferroptosis, parthanatos, and pyroptosis may be occurring. The role of M1 microglial cells in initiating motor neuron destruction and M₂ cells in ameliorating destruction is unknown. Use of TBI and bone marrow transplantation might be possibly in other neurodegenerative diseases including: multiple sclerosis, frontotemporal dementia, and the multiple etiologies of sporadic ALS such as that following trauma.

SOD1^{G93A} mice receiving bone marrow transplant at day 60 before onset of signs of paralysis fared much better, but translation of these data to the clinic is not practical since diagnosis of ALS is made after the onset of symptoms and weakness. Our mouse data do suggest that the course of degeneration might be further interrupted by earlier bone marrow transplantation. The mechanisms for blood–brain




Figure 7. Radiation resistance of fresh marrow hematopoietic progenitor cells from superoxide dismutase-1 (SOD1)^{G93A} mice (n=3) (A) and bone marrow stromal cell lines from SOD1^{G93A} and SOD1 transgenic (B6) mice (B). Radiation survival curves were carried out as described in the Materials and Method. The data are a composite of data from 3-6 experiments.

barrier permeability in ALS, and its restoration by marrow transplant is at present not known.

The unexpected prolongation of LTBMHC hematopoiesis and the radioresistance of cell lines and fresh marrow from SOD1^{G93A} mice appears to be unrelated to motor neuron defects in these same mice. SOD1^{G93A} mice with four copies of the mutant gene compared to copies of the normal human gene showed no increase in duration of hematopoiesis in LTBMHCs, and cell lines were not radioresistant. The defective SOD1 may have caused changes in the bone marrow resulting in the up-regulation of SOD2, glutathione, or glutathione peroxidase-4 that may explain the differences in bone marrow biology in SOD1^{G93A} mice.

TBI combined with bone marrow transplantation is a potentially valuable therapeutic option for treating ALS. The subset of donor bone marrow cells (hematopoietic stem cells, committed granulocyte/macrophage progenitors, or mesenchymal stem cells) required to provide the therapeutic effect must also be determined (59, 60, 63, 64, 79-82).

Lrcc33^{-/-} mice lack central nervous system vascular abnormalities. These mice have a different transformed growth factor-beta, develop ascending paraparesis and death attributable to microglia activation. Recently it was




Figure 8. Stages of spinal cord paralysis in superoxide dismutase-1 (SOD1)^{G93A} mice. Marrow-transplanted and control SOD1^{G93A} mice were sacrificed when developing amyotrophic lateral sclerosis with neurological score (NS) 0, 1, 2 or 3 paralysis (n=3). The spine was removed from euthanized mice, and the spinal cord was flushed from the vertebral column, fixed for 2 h in 2% paraformaldehyde, and then stored in 30% sucrose for 24 h. The number of neurons in sections at each stage of paralysis was normalized by dividing the number of neurons by the number of nuclei.

demonstrated that transplantation of donor bone marrow from a wild-type mouse ameliorated these abnormalities due to the migration of wild type bone marrow origin microglia from the bone marrow (75). These data are a further indication in another model system that microglia arising from transplanted marrow can ameliorate a spinal cord degenerative disorder. Microglia appears to be a central component of the pathophysiology of ALS, as well as other neurodegenerative disorders (76), and bone marrow transplants may provide a cellular mechanism to correct such defects.

Supplemental Data

All supplemental data may be found at <https://pitt.box.com/s/ajzj6ov2cw9ttrbnsfnfz3p7d7so7mob>

Conflicts of Interest

The Authors have no conflicts of interest in regard to this study.

Authors' Contributions

Michael Epperly, Ph.D., supervised all experiments. Joel Greenberger, M.D., planned and reviewed data for every experiment. Stephanie Theriozier scored radiation survival assay, and slides for immunostaining. Lora Rigatti, VMD, reviewed histopathology of spinal cords. Xichen Zhang, MD, carried out analysis of genotyping. Donna Shields carried out long-term bone marrow cultures. Wen Hou carried aided in staining of slides. Renee Fisher carried out all animal breeding and genotyping, and, also, observed animals daily and documented stages of paralysis. Simon Watkins, Ph.D., prepared fluorescent images. Andrew Henderson carried out the analysis of the numbers of neurons in spinal cords. Darcy Francicola carried out the assays for antioxidants and radiation survival curves of cell lines and fresh marrow. Christopher Donnelly, Ph.D., carried out review of all data, planned experiments in collaboration with radiation oncologists, and served as the neurosciences contributor for analysis and interpretation of all data. Hong Wang, Ph.D., completed all the biostatistics for the article. Hulya Bayir, M.D., analyzed histopathology and stages of paralysis. Peter Wipf, Ph.D., prepared drugs tested in Figure 9.

Acknowledgements

This study was supported by NIH U19-A1068021 grant. Presented at the American Society of Therapeutic Radiology and Oncology (ASTRO) Annual Meeting, October, 2018, San Antonio, TX, USA.

References

- Haverkamp LJ, Appel V and Appel SH: Natural history of amyotrophic lateral sclerosis in a database population. Validation of a scoring system and a model for survival prediction. *Brain* 118(pt3): 707-719, 1995. PMID: 7600088.
- Boillee S, Vande Velde C and Cleveland DW: ALS: a disease of motor neurons and their nonneuronal neighbors. *Neuron* 52(1): 39-59, 2006. PMID: 17015226.
- Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, Donaldson D, Goto J, O'Regan JP and Deng HX: Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* 364(6435): 362, 1993. PMID: 8446170. DOI: 10.1038/362059a0
- Sreedharan J, Blair IP, Tripathi VB, Hu X, Vance C, Rogelj B, Ackerley S, Durnall JC, Williams KL, Buratti E, Baralle F, de Bellerocche J, Mitchell JD, Leigh PN, Al-Chalabi A, Miller CC, Nicholson G and Shaw CE: TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis. *Science* 319(5870): 1668-1672, 2008. PMID: 18309045. DOI: 10.1126/science.1154584
- Vance C, Rogelj B, Hortobagyi T, De Vox KJ, Nishimura AL, Sreedharan J, Hu X, Smith B, Ruddy D, Wright P, Ganesalingam J, Williams KL, Tripathi V, Al-Saraj S, Al-Chalabi A, Leigh PN, Blair IP, Nicholson G, de Bellerocche J, Gallo JM, Miller CC and Shaw CE: Mutations in *FUS*, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science* 323(5918): 1208-1211, 2009. PMID: 19251628. DOI: 10.1126/science.1165942
- Kwiatkowski, Jr TJ, Bosco DA, Leclerc AL, Tamrazian E, Vanderburg CR, Russ C, Davis A, Gilchrist J, Kasarskis EJ, Munsat T, Valdmanis P, Rouleau GA, Hosler BA, Cortelli P, de Jong PJ, Yoshinaga Y, Haines JL, Pericak-Vance MA, Yan J, Ticozzi N, Siddique T, McKenna-Yasek D, Sapp PC, Horvitz HR, Landers JE and Brown RH Jr.: Mutations in the *FUS/ALS* gene on chromosome 16 cause familial amyotrophic lateral sclerosis. *Science* 323(5918): 1205-1208, 2009. PMID: 19251627. DOI: 10.1126/science.1166066
- Fecto F, Yan J, Vemula SP, Liu E, Yang Y, Chen W, Zheng JG, Shi Y, Siddique N, Arrat H, Donkervoort S, Ajroud-Driss S, Sufit RL, Heller SL, Deng HX and Siddique T: *SQSTM1* mutations in familial and sporadic amyotrophic lateral sclerosis. *Arch Neurol* 68(11): 1440-1446, 2011. PMID: 22084127. DOI: 10.1001/archneurol.2011.250
- Turner, MR, Cagnin A, Turkheimer FE, Miller CCJ, Shaw CE, Brooks DJ, Leigh PN and Banati RB: Evidence of wide spread cerebral microglia activation in amyotrophic lateral sclerosis: An [¹¹C] @-PK11195 positron emission tomography study. *Neurobiol Dis* 15: 601-609, 2004. PMID: 15056468.
- Lall D and Baloh RH: Microglia and C9orf72 in neuroinflammation and ALS and frontotemporal dementia. *J Clin Invest* 127(9): 3250-3258, 2017. PMID: 28737506. DOI: 10.1172/JCI90607
- Matcovitch-Natan O, Winter DR, Giladi A, Vargas Aguilar S, Spinrad A, Sarrazin S, Ben-Yehuda H, David E, Zelada Gonzalez F, Perrin P, Keren-Shaul H, Gury M, Lara-Astaiso D, Thaiss CA, Cohen M, Bahar Halpern K, Baruch K, Deczkowska A, Lorenzo-Vivas E, Itzkovitz S, Elinav E, Sieweke MH, Schwartz M and Amit I: Microglia development follows a stepwise program to regulate brain homeostasis. *Science* 353(6301): aad8670, 2016. PMID: 27338705. DOI: 10.1126/science.aad8670
- Town T, Nikolic V and Tan J: The microglial "activation" continuum: from innate to adaptive responses. *J Neuroinflammation* 2: 24, 2005. PMID: 16259628. DOI: 10.1186/1742-2094-2-24
- Ginhoux F, Lim S, Hoeffel G, Low D and Huber T: Origin and differentiation of microglia. *Front Cell Neurosci* 7: 45, 2013. PMID: 23616747. DOI: 10.3389/fncel.2013.00045
- London A, Cohen M and Schwartz M: Microglia and monocyte-derived macrophages: functionally distinct populations that act in concert in CNS plasticity and repair. *Front Cell Neurosci* 7: 34, 2013. PMID: 23596391. DOI: 10.3389/fncel.2013.00034

- 14 Popovich PG and Hickey WF: Bone marrow chimeric rats reveal the unique distribution of resident and recruited macrophages in the contused rat spinal cord. *J Neuropathol Exp Neurol* 60(7): 676-685, 2001. PMID: 11444796.
- 15 Ajami B, Bennett JL, Krieger C, McNagny KM and Rossi FM: Infiltrating monocytes trigger EAE progression, but do not contribute to the resident microglia pool. *Nat Neurosci* 14(9): 1142-1149, 2011. PMID: 21804537. DOI: 10.1038/nn.2887
- 16 Yamanaka K, Boillee S, Roberts EA, Garcia ML, McAlonis-Downes M, Mikse OR, Cleveland DW and Goldstein LS: Mutant SOD1 in cell types other than motor neurons and oligodendrocytes accelerates onset of disease in SOD1^{G93A} mice. *Proc Natl Acad Sci USA* 105(21): 7594-7599, 2008. PMID: 18492803. DOI: 10.1073/pnas.0802556105
- 17 Jaarsma D, Teuling E, Haasdijk ED, DeZeeuw CI and Hoogenraad CC: Neuron-specific expression of mutant superoxide dismutase is sufficient to induce amyotrophic lateral sclerosis in transgenic mice. *J Neurosci* 28(9): 2075-2088, 2008. PMID: 18305242. DOI: 10.1523/JNEUROSCI.5258-07.2008
- 18 Clement AM, Nguyen MD, Roberts EA, Garcia ML, Boillee S, Rule M, McMahon AP, Doucette W, Siwek D, Ferrante RJ, Brown RH Jr., Julien JP, Goldstein LS and Cleveland DW: Wild-type nonneuronal cells extend survival of SOD1 mutant motor neurons in SOD1^{G93A} mice. *Science* 302(5642): 113-117, 2003. PMID: 14526083. DOI: 10.1126/science.1086071
- 19 Zhao W, Xie W, Le W, Beers DR, He Y, Henkel JS, Simpson EP, Yen AA, Xiao Q and Appel SH: Activated microglia initiate motor neuron injury by a nitric oxide and glutamate-mediated mechanism. *J Neuropathol Exp Neurol* 63(9): 964-977, 2004. PMID: 15453095.
- 20 Xiao Q, Zhao W, Beers DR, Yen AA, Xie W, Henkel JS and Appel SH: Mutant SOD (G93A) microglia are more neurotoxic relative to wild-type microglia. *J Neurochem* 102(6): 2008-2019, 2007. PMID: 17555556. DOI: 10.1111/j.1471-4159.2007.04677.x
- 21 Weydt P, Yuen EC, Ransom BR and Moller T: Increased cytotoxic potential of microglia from ALS-transgenic mice. *Glia* 48(2): 179-182, 2004. PMID: 15378658. DOI: 10.1002/glia.20062
- 22 Frakes AE, Ferraiuolo L, Haidet-Phillips AM, Schmeizer L, Braun L, Miranda CJ, Ladner KJ, Bevan AK, Foust KD, Godbout JP, Popovich PG, Guttridge DC and Kaspar BK: Microglia induce motor neuron death via the classical NF- κ B pathway in amyotrophic lateral sclerosis. *Neuron* 81(5): 1009-1023, 2014. PMID: 24607225. DOI: 10.1016/j.neuron.2014.01.013
- 23 Alexianu ME, Kozovska M and Appel SH: Immune reactivity in a mouse model of familial ALS correlates with disease progression. *Neurology* 57(7): 1282-1289, 2001. PMID: 11591849.
- 24 Zhao W, Beers DR, Henkel JS, Zhang W, Urushitani M, Julien JP and Appel SH: Extracellular mutant SOD1 induces microglial-mediated motoneuron injury. *Glia* 58(2): 231-243, 2010. PMID: 19672969. DOI: 10.1002/glia.20919
- 25 Harraz MM, Marden JJ, Zhou W, Zhang Y, Williams A, Sharov VS, Nelson K, Luo M, Paulson H, Schoneich C and Engelhardt JF: SOD1 mutations disrupt redox-sensitive Rac regulation of NADPH oxidase in a familial ALS model. *J Clin Invest* 118(2): 659-670, 2008. PMID: 18219391. DOI: 10.1172/JCI34060
- 26 Hatzipetros T, Kidd JD, Moreno AJ, Thompson K, Gill A and Vieira FG: A quick phenotype neurological scoring system for evaluating disease progression in the SOD1-G93A mouse model of ALS. *J Visual Exp Oct* 6(104): e53257, 2015. PMID: 26485052. DOI: 10.3791/53257
- 27 Liao B, Zhao W, Beers DR, Henkel JS and Appel SH: Transformation from a neuroprotective to a neurotoxic microglial phenotype in a mouse model of ALS. *Exp Neurol* 237(1): 147-152, 2012. PMID: 22735487. DOI: 10.1016/j.expneurol.2012.06.011
- 28 Brettschneider J, Toledo JB, Van Deerlin VM, Elman L, McCluskey L, Lee VM and Trojanowski JQ: Microglial activation correlates with disease progression and upper motor neuron clinical symptoms in amyotrophic lateral sclerosis. *PLoSOne* 7(6): e39216, 2012. PMID: 22720079. DOI: 10.1371/journal.pone.0039216
- 29 Tarhini Ahmad A, Belani Chandra, Luketich James D, Ramalingam Suresh S, Argiris Athanassios, Gooding William, Pennathur A, Petro D, Kane K, Liggitt D, Championsmith T, Zhang X, Epperly MW and Greenberger JS: A phase I study of concurrent chemotherapy (paclitaxel and carboplatin) and thoracic radiotherapy with swallowed manganese superoxide dismutase (MnSOD) plasmid liposome (PL) protection in patients with locally advanced stage III non-small cell lung cancer. *Hum Gene Ther* 22(3): 336-343, 2011. PMID: 20873987. DOI: 10.1089/hum.2010.078
- 30 Ferraiuolo L, Meyer K, Sherwood TW, Vick J, Likhite S, Frakes A, Miranda CJ, Braun L, Heath PR, Pineda R, Beattie CE, Shaw PJ, Askwith CC, McTigue D and Kaspar BK: Oligodendrocytes contribute to motor neuron death in ALS via SOD1-dependent mechanism. *Proc Natl Acad Sci USA* 113(42): E6946-E6505, 2016. PMID: 27688759. DOI: 10.1073/pnas.1607496113
- 31 Buttgereit A, Lelios I, Yu X, Vrohings M, Krakoski NR, Gautier EL, Nishinakamura R, Becher B and Greter M: Sall1 is a transcriptional regulator defining microglia identity and function. *Nat Immunol* 17: 1397-1406, 2016. PMID: 27776109. DOI: 10.1038/ni.3585
- 32 Taylor JP, Brown, Jr. RH and Cleveland DW: Decoding ALS: from genes to mechanism. *Nature* 539: 197-206, 2016. PMID: 27830784. DOI: 10.1038/nature20413
- 33 Falkner S, Grade S, Dimou L, Conzelmann K-K, Bonhoeffer T, Gotz M and Hubener M: Transplanted embryonic neurons integrate into adult neocortical circuits. *Nature* 539: 248-253, 2016. PMID: 27783592. DOI: 10.1038/nature20113
- 34 Sunico CR, Sultan A, Nakamura T, Dolatabadi N, Parker J, Shan B, Han X, Yates JR 3rd, Maslah E, Ambasudhan R, Nakanishi N and Lipton SA: Role of sulfiredoxin as a peroxiredoxin-2 denitrosylase in human iPSC-derived dopaminergic neurons. *Proc Natl Acad Sci USA* 113: E7564-E7571, 2016. PMID: 27821734. DOI: 10.1073/pnas.1608784113
- 35 Iaccarino HF, Singer AC, Martorell AJ, Rudenko A, Gao F, Gillingham TZ, Mathys H, Seo J, Kritskiy O, Abdurrob F, Adaikkan C, Canter RG, Rueda R, Brown EN, Boyden ES and Tsai LH: Gamma frequency entrainment attenuates amyloid load and modifies microglia. *Nature* 540: 230-235, 2016. PMID: 27929004. DOI: 10.1038/nature20587
- 36 Lecuyer M-A, Saint-Laurent O, Bourbonniere L, Larouche S, Larochelle C, Michel L, Charabati M, Abadier M, Zandee S, Haghayegh Jahromi N, Gowing E, Pittet C, Lyck R, Engelhardt B and Prat A: Dual role of ALCAM in neuroinflammation and blood-brain barrier homeostasis. *Proc Natl Acad Sci USA* 114: E524-E533, 2017. PMID: 28069965. DOI: 10.1073/pnas.1615336114
- 37 Engelhardt B, Vajkoczy P and Weller RO: The movers and shapers in immune privilege of the CNS. *Nat Immunol* 18(2): 123-133, 2017. PMID: 28092374. DOI: 10.1038/ni.3666

- 38 Liddel SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, Bennett ML, Munch AE, Chung WS, Peterson TC, Wilton DK, Frouin A, Napier BA, Panicker N, Kumar M, Buckwalter MS, Rowitch DH, Dawson VL, Dawson TM, Stevens B and Barres BA: Neurotoxic reactive astrocytes are induced by activated microglia. *Nature* 541: 481-487, 2017. PMID: 28099414. DOI: 10.1038/nature21029
- 39 Velasco S, Ibrahim MM, Kakumanu A, Garipler G, Aydin B, Al-Sayegh MA, Hirsekorn A, Abdul-Rahman F, Satija R, Ohler U, Mahony S and Mazzone EO: A multi-step transcriptional and chromatin state cascade underlies motor neuron programming from embryonic stem cells. *Cell Stem Cell* 20: 205-217, 2017. PMID: 27939218. DOI: 10.1016/j.stem.2016.11.006
- 40 Ehrlich M, Mozafari S, Glatza M, Starost L, Velychko S, Hallmann A-L, Cui QL, Schambach A, Kim KP, Bachelin C, Marteyn A, Hargus G, Johnson RM, Antel J, Sterneckert J, Zaehres H, Scholer HR, Baron-Van Evercooren A and Kuhlmann T: Rapid and efficient generation of oligodendrocytes from human induced pluripotent stem cells using transcription factors. *Proc Natl Acad Sci USA* 114: E2243-E2252, 2017. PMID: 28246330. DOI: 10.1073/pnas.1614412114
- 41 Chang J, Mancuso MR, Maier C, Liang X, Yuki K, Yang L, Kwong JW, Wang J, Rao V, Vallon M, Kosinski C, Zhang JJ, Mah AT, Xu L, Li L, Gholamin S, Reyes TF, Li R, Kuhnert F, Han X, Yuan J, Chiou SH, Brettman AD, Daly L, Corney DC, Cheshier SH, Shortliffe LD, Wu X, Snyder M, Chan P, Giffard RG, Chang HY, Andreasson K and Kuo CJ: Gpr124 is essential for blood-brain barrier integrity in central nervous system disease. *Nat Med* 23(4): 450-460, 2017. PMID: 28288111. DOI: 10.1038/nm.4309
- 42 Haidet-Phillips AM, Doreswamy A, Gross SK, Tang X, Campanelli JT and Maragakis NJ: Human glial progenitor engraftment and gene expression is independent of the ALS environment. *Exp Neurol* 264: 343-350, 2015. PMID: 25523812. DOI: 10.1016/j.expneurol.2014.12.011
- 43 Haidet-Phillips AM and Maragakis NJ: Neural and glial progenitor transplantation as a neuroprotective strategy for amyotrophic lateral sclerosis (ALS). *Brain Res* 1628: 343-350, 2015. PMID: 26187754. DOI: 10.1016/j.brainres.2015.06.035
- 44 Kondo T, Funayama M, Tsukita K, Hotta A, Yasuda A, Nori S, Kaneko S, Nakamura M, Takahashi R, Okano H, Yamanaka S and Inoue H: Focal transplantation of human iPSC-derived glial-rich neural progenitors improves lifespan of SOD1^{G93A} mice. *Stem Cell Reports* 3: 242-249, 2014. PMID: 25254338. DOI: 10.1016/j.stemcr.2014.05.017
- 45 Chen H, Qian K, Chen W, Hu B, Blackbourn, IV LW, Du Z, Ma L, Liu H, Knobel KM, Ayala M and Zhang SC: Human-derived neural progenitors functionally replace astrocytes in adult mice. *J Clin Invest* 125(3): 1033-1043, 2015. PMID: 25642771. DOI: 10.1172/JCI69097
- 46 Nicaise C, Mitrecic D, Fahnrikar A and Lepore AC: Transplantation of stem cell-derived astrocytes for the treatment of amyotrophic lateral sclerosis and spinal cord injury. *World J Stem Cells* 7(2): 380-398, 2015. PMID: 25815122. DOI: 10.4252/wjsc.v7.12.380
- 47 Winkler EA, Sengillo JD, Sagare AP, Zhao Z, Ma Q, Zuniga E, Wang Y, Zhong Z, Sullivan JS, Griffin JH, Cleveland DW and Zlokovic BV: Blood-spinal cord barrier disruption contributes to early motor-neuron degeneration in ALS-model mice. *Proc Natl Acad Sci USA* 111(11): E1035-E1042, 2014. PMID: 24591593. DOI: 10.1073/pnas.1401595111
- 48 Haeusler AR, Donnelly CJ and Rothstein JD: The expanding biology of the C9orf72 nucleotide repeat expansion in neurodegenerative disease. *Nature Rev Neurosci* 17(6): 383-395, 2016. PMID: 27150398. DOI: 10.1038/nrn.2016.38
- 49 Hill-Kayser CE, Plastaras JP, Tochner Z and Glatstein E: TBI during BM and SCT: review of the past, discussion of the present, and consideration of future directions. *Bone Marrow Transplant* 46: 475-484, 2011. PMID: 21113184. DOI: 10.1038/bmt.2010.280
- 50 Schneiderman J, Thormann K and Kletzel M: Correction of enzyme levels with allogeneic hematopoietic progenitor cell transplantation in Niemann-Pick type B. *Pediatr Blood Cancer*: 49(7): 987-989, 2007. PMID: 17635007 DOI: 10.1002/pbc.21300
- 51 Appel SH, Engelhardt JI, Henkel JS, Siklos L, Beers DR, Yen AA, Simpson EP, Luo Y, Carrum G, Heslop HE, Brenner MK and Popat U: Hematopoietic stem cell transplantation in patients with sporadic amyotrophic lateral sclerosis. *Neurology* 71(17): 1326-1334, 2008. PMID: 18936424. DOI: 10.1212/01.wnl.0000327668.43541.22
- 52 Ohnishi S, Ito H, Suzuki Y, Adachi Y, Wate R, Zhang J, Nakano S, Kusaka H and Ikehara S: Intra-bone marrow-bone marrow transplantation slows disease progression and prolongs survival in G93A mutant SOD1 transgenic mice, an animal model mouse for amyotrophic lateral sclerosis. *Brain Res* 1296: 216-224, 2009. PMID: 19686706. DOI: 10.1016/j.brainres.2009.08.012
- 53 Van der Kogel AJ: Dose-volume effects in the spinal cord. *Radiother Oncol* 29(2): 105-109, 1993. PMID: 8310135.
- 54 Jiang Y, Jahagirdar BN, Reinhardt RL, Schwartz RE, Keene CD, Ortiz-Gonzalez XR, Reyes M, Lenvik T, Lund T, Blackstad M, Du J, Aldrich S, Lisberg A, Low WC, Largaespada DA and Verfaillie CM: Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature* 418: 41-49, 2002. PMID: 12077603. DOI: 10.1038/nature00870
- 55 Morikawa S, Mabuchi Y, Kubota Y, Nagai Y, Niibe K, Hiratsu E, Suzuki S, Miyauchi-Hara C, Nagoshi N, Sunabori T, Shimmura S, Miyawaki A, Nakagawa T, Suda T, Okano H and Matsuzaki Y: Prospective identification, isolation, and systemic transplantation of multipotent mesenchymal stem cells in murine bone marrow. *J Exp Med* 206(11): 2483-2496, 2009. PMID: 19841085. DOI: 10.1084/jem.20091046
- 56 Yu H, Shen H, Yuan Y, XuFeng R, Hu X, Garrison SP, Zhang L, Yu J, Zambetti GP and Cheng T: Deletion of Puma protects hematopoietic stem cells and confers long-term survival in response to high-dose γ -irradiation. *Blood* 115: 3472-3480, 2010. PMID: 20177048. DOI: 10.1182/blood-2009-10-248278
- 57 Gowing G, Philips T, Van Wijmeersch B, Audet JN, Dewil M, Van Den Bosch L, Billiau AD, Robberecht W and Julien JP: Ablation of proliferating microglia does not affect motor neuron degeneration in amyotrophic lateral sclerosis caused by mutant superoxide dismutase. *J Neurosci* 28: 10234-10244, 2008. PMID: 18842883. DOI: 10.1523/JNEUROSCI.3494-08.2008
- 58 Lino MM, Schneider C and Caroni P: Accumulation of SOD1 mutants in postnatal motoneurons does not cause motoneuron pathology or motoneuron disease. *J Neurosci* 22: 4825-4832, 2002. PMID: 12077179.
- 59 Willis J, Epperly MW, Fisher R, Zhang X, Shields D, Hou W, Wang H, Li S, Wipf P, Parmar K, Guinan E, Steinman J and Greenberger JS: Amelioration of head and neck radiation-induced mucositis and distant marrow suppression in *Fanca*^{-/-} and *Fancg*^{-/-} mice by intraoral administrations of GS-nitroxide (JP4-039). *Radiat Res* 189: 560-578, 2018. PMID: 29584588. DOI: 10.1667/RR14878.1

- 60 Epperly MW, Sikora CA, Defilippi S, Gretton JE and Greenberger JS: Bone marrow origin of myofibroblasts in irradiation pulmonary fibrosis. *Am J Resp Molecular Cell Biol.* 29: 213-224, 2003. PMID: 12649121. DOI: 10.1165/rcmb.2002-0069OC
- 61 Kalash R, Epperly MW, Goff J, Dixon T, Sprachman MM, Zhang X, Shields D, Cao S, Wipf P, Francicola D, Berhane H and Greenberger JS: Amelioration of irradiation pulmonary fibrosis by a water-soluble bi-functional sulfoxide radiation mitigator (MMS350). *Radiat Res* 180: 474-490, 2013. PMID: 24125487. DOI: 10.1667/RR15199.1
- 62 Sivananthan A, Shields D, Fisher R, Hou W, Zhang X, Francicola D, Epperly MW, Wipf P and Greenberger JS: Continuous 1 year oral administration of the radiation mitigator, MMS350, after total body irradiation restores bone marrow stromal cell proliferative capacity and reduces age-related senescence in Fanconi Anemia (*Fanca*^{-/-}) mice. *Radiat Res* 191: 139-143, 2019. PMID: 30499383. DOI: 10.1667/RR15199.1
- 63 Berhane H, Epperly MW, Goff J, Kalash R, Cao S, Francicola D, Zhang X, Shields D, Houghton F, Wang H, Wipf P, Parmar K and Greenberger JS: Radiologic differences between bone marrow stromal and hematopoietic progenitor cell lines from Fanconi Anemia (*Fancd2*^{-/-}) mice. *Radiat Res* 181(1): 76-89, 2014. PMID: 24397476. DOI: 10.1667/RR13405.1
- 64 Shinde A, Berhane H, Rhieu BH, Kalash R, Xu K, Goff J, Epperly MW, Francicola D, Zhang X, Dixon T, Shields D, Wang H, Wipf P, Parmar K, Guinan E, Kagan V, Tyurin V, Ferris RL, Zhang X, Li S and Greenberger JS: Intraoral mitochondrial-targeted GS-Nitroxide, JP4-039, radioprotects normal tissue in tumor-bearing radiosensitive *Fancd2*^{-/-} (C57BL/6) mice. *Radiat Res* 185: 134-150, 2016. PMID: 26789701. DOI: 10.1667/RR14035.1
- 65 Watson AM, Watkins S, Sivananthan A, Thermoziar S and Greenberger JS: Chapter XXIV: *In situ* correlations of radiation damage: from the single molecule to the whole animal. *In: Radiobiology Methods Web-Based Course*, Pittsburgh, PA, USA, 2019. Available from: <https://pitt.app.box.com/v/CMCR-Virtual-Publications-2018?sortColumn=name&sortDirection=ASC>
- 66 Anklesaria P, Kase KR, Glowacki J, Holland CH, Sakakeeny MA, Wright JH, FitzGerald TJ, Lee CY and Greenberger JS: Engraftment of a clonal bone marrow stromal cell line *in vivo* stimulates hematopoietic recovery from total body irradiation. *Proc Natl Acad Sci USA* 84: 7681-7685, 1987. PMID: 2890167.
- 67 Greenberger JS, FitzGerald TJ, Kleasen V, Anklesaria P, Bushnell D, Kase K and Sakakeeny MA: Alteration in hematopoietic stem cell seeding and proliferation by low-dose-rate irradiation of bone marrow stromal cells *in vitro*. *Int J Radiat Oncol Biol Phys* 14: 85-94, 1988. PMID: 2891657.
- 68 Epperly MW, Epstein CJ, Travis EL and Greenberger JS: Decreased pulmonary radiation resistance of manganese superoxide dismutase (MnSOD)-deficient mice is corrected by human manganese superoxide dismutase-plasmid/liposome (SOD2-PL) intratracheal gene therapy. *Radiat Res* 154(4): 365-374, 2000. PMID: 11023599.
- 69 Epperly MW, Bernarding M, Gretton J, Jefferson M, Nie S and Greenberger JS: Overexpression of the transgene for manganese superoxide dismutase (MnSOD) in 32D cl 3 cells prevent apoptosis induction by TNF- α , IL-3 withdrawal, and ionizing irradiation. *Exp Hematol* 31(6): 465-474, 2003.
- 70 Epperly MW, Goff JP, Sikora CA, Shields DS and Greenberger JS: Bone marrow origin of cells with capacity for homing and differentiation to esophageal squamous epithelium. *Radiat Res* 162: 233-240, 2004. PMID: 15333000.
- 71 Bernard ME, Kim H, Rwigema J-C, Epperly MW, Kelley EE, Murdoch GH, Dixon T, Wang H and Greenberger JS: Role of the esophageal vagus neural pathway in ionizing irradiation-induced seizures in nitric oxide synthase-1 homologous recombination negative *NOS1*^{-/-} mice. *In Vivo* 25(6): 861-870, 2011.
- 72 Vintersten K, Monetti C, Gertsenstein M, Zhang P, Laszlo L, Biechele S and Nagy A: Mouse in red: Red fluorescent protein expression in mouse ES cells, embryos, and adult animals. *Genesis* 40(4): 241-246, 2004. PMID: 15593332. DOI: 10.1002/gene.20095
- 73 Butovsky O and Weiner HL: Microglial signatures and their role in health and disease. *Nat Rev Neurosci* 19(62): 622-634, 2018. PMID: 30206328. DOI: 10.1038/s41583-018-0057-5
- 74 Heneka MT, McManus RM and Latz E: Inflammasome signalling in brain function and neurodegenerative disease. *Nat Rev Neurosci* 19: 610-621, 2018. PMID: 30206330. DOI: 10.1038/s41583-018-0055-7
- 75 Johann S, Heitzer M, Kanagaratnam M, Goswami A, Rizo T, Weis J, Troost D and Beyer C: NLRP3 inflammasome is expressed by astrocytes in the SOD1 mouse model of ALS and in human sporadic ALS patients. *Glia* 63(12): 2260-2273, 2015. PMID: 26200799. DOI: 10.1002/glia.22891
- 76 Meissner F, Molawi K and Zychlinsky A: Mutant superoxide dismutase 1-induced IL-1 beta accelerates ALS pathogenesis. *Proc Natl Acad Sci USA* 107: 13046-13050, 2010. PMID: 20616033. DOI: 10.1073/pnas.1002396107
- 77 Bellezza I, Grottelli S, Costanzi E, Scarpelli P, Pigna E, Morozzi G, Mezzasoma L, Peirce MJ, Moresi V, Adamo S and Minelli A: Peroxynitrite activates the NLRP3 inflammasome cascade in SOD1 (G93A) mouse model of amyotrophic lateral sclerosis. *Mol Neurobiol* 55(3): 2350-2361, 2018. PMID: 28357805. DOI: 10.1007/s12035-017-0502-x
- 78 Italiani P, Carlesi C, Giungato P, Puxeddu I, Borroni B, Bossu P, Migliorini P, Siciliano G and Boraschi D: Evaluating the levels of interleukin-1 family cytokines in sporadic amyotrophic lateral sclerosis. *J Neuroinflammation* 11: 94, 2014. PMID: 24884937. DOI: 10.1186/1742-2094-11-94
- 79 Qin Y, Garrison BS, Ma W, Wang R, Jiang A, Li J, Mistry M, Bronson RT, Santoro D, Franco C, Robinton DA, Stevens B, Rossi DJ, Lu C and Springer TA: A milieu molecule for TGF- β required for microglia function in the nervous system. *Cell* 174: 156-171, 2018. PMID: 29909984. DOI: 10.1016/j.cell.2018.05.027
- 80 Deczkowska A, Keren-Shaul H, Weiner A, Colonna M, Schwartz M and Amit I: Disease-associated microglia: a universal immune sensor of neurodegeneration. *Cell* 173: 1073-1083, 2018. PMID: 29775591. DOI: 10.1016/j.cell.2018.05.003
- 81 Steinman J, Epperly M, Hou W, Willis J, Wang H, Fisher R, Liu B, Bahar I, McCaw T, Kagan V, Bayir H, Yu J, Wipf P, Li S, Huq MS and Greenberger JS: Improved total-body irradiation survival by delivery of two radiation mitigators that target distinct cell death pathways improved total-body irradiation survival by delivery of two radiation mitigators that target distinct cell death pathways. *Radiat Res* 189(1): 68-83, 2018. PMID: 29140165. DOI: 10.1667/RR14787.1
- 82 Berhane H, Shinde A, Kalash R, Xu K, Epperly MW, Goff J, Francicola D, Zhang X, Dixon T, Shields D, Wang H, Wipf P, Li S, Gao X and Greenberger JS: Amelioration of irradiation induced oral cavity mucositis and distant bone marrow suppression in *Fancd2*^{-/-} (FVB/N) mice by intraoral JP4-039/F15. *Radiat Res* 182: 35-49, 2014. PMID: 24932534. DOI: 10.1667/RR13633.1

Received February 18, 2019

Revised March 29, 2019

Accepted April 2, 2019