

Editorial Board

- N.J. AGNANTIS, *Department of Pathology, University of Ioannina, Ioannina, Greece*
D. ANDERSON, *Department of Biomedical Sciences, University of Bradford, Bradford, West Yorkshire, UK*
V. BARAK, *Department of Oncology, Hadassah University Hospital, Jerusalem, Israel*
R.J.B. BASSLEER, *Department of Histology, Université de Liège, Liège, Belgium*
Y. BECKER, *Department of Molecular Biology, Hebrew University of Jerusalem, Jerusalem, Israel*
K. BEIER, *Department of Histology, University of Basel, Basel, Switzerland*
S. BEN-EFRAIM, *Department of Human Microbiology, University of Tel-Aviv, Tel-Aviv, Israel*
J. BEUTH, *Institute of Scientific Evaluation of Naturopathy, University of Köln, Köln, Germany*
R. BJERKVIG, *Department of Anatomy and Cell Biology, University of Bergen, Bergen, Norway*
B. BODEY, *Department of Pathology, University of Southern California, Reseda, CA, USA*
K. W. BRUNSON, *Department of Molecular Biology and Immunology, University of North Texas, Fort Worth, TX, USA*
P. CHANDRA, *Department of Molecular Biology, Frankfurt University, Frankfurt am Main, Germany*
J.-G. CHUNG, *Department of Medicine, China Medical College, Taichung, Taiwan*
L.A. COHEN, *Institute for Cancer Prevention, Valhalla, NY, USA*
A.I. CONSTANTINOU, *Department of Biological Sciences, University of Cyprus, Nicosia, Cyprus*
J.G. CORY, *Department of Biochemistry and Molecular Biology, East Carolina University, Greenville, NC, USA*
T. DALIANIS, *Department of Pathology-Oncology, Karolinska Institute, Stockholm, Sweden*
I. DAMJANOV, *Department of Pathology, The University of Kansas, Kansas City, KS, USA*
G. DELICONSTANTINOS, *Department of Experimental Physiology, University of Athens Medical School, Athens, Greece*
D.T. DENHARDT, *Department of Cell Biology and Neurosciences, Rutgers University, Piscataway, NJ, USA*
W. DEN OTTER, *Department of Biochemistry, University of Utrecht, Utrecht, The Netherlands*
L. DE RIDDER, *Department of Anatomy, Embryology and Histology, Universiteit Gent, Gent, Belgium*
E.P. DIAMANDIS, *Department of Pathology and Laboratory Medicine, University of Toronto, Toronto, Ont., Canada*
J.F. DORÉ, *Centre Léon Berard, Lyon, France*
E. EIGENBRODT, *Institute for Biochemistry and Endocrinology, Justus-Liebig-University of Giessen, Giessen, Germany*
I. EMBER, *Department of Public Health, University of Pécs Medical School, Pécs, Hungary*
W. ENGSTRÖM, *Department of Molecular Biosciences, Swedish University of Agricultural Sciences, Uppsala, Sweden*
M. ESKELINEN, *Department of Surgery, University Hospital of Kuopio, Kuopio, Finland*
J.A. FERNANDEZ-POL, *Metalloproteomics, LLC, Chesterfield, MO, USA*
S. FERRONE, *Department of Immunology, Roswell Park Cancer Institute, Buffalo, NY, USA*
P.B. FISHER, *Department of Pathology, Neurosurgery and Urology, Columbia University, New York, NY, USA*
I. FREITAS, *Dipartimento di Biologia Animale, University of Pavia, Pavia, Italy*
R.E. FRIEDRICH, *Department of Oral and Maxillofacial Surgery, Eppendorf University Hospital, Hamburg, Germany*
R. GANAPATHI, *Taussig Cancer Center, The Cleveland Clinic Foundation, Cleveland, OH, USA*
D.H. GILDEN, *Department of Neurology, University of Colorado, Denver, CO, USA*
G. GITSCH, *Department of Gynecology and Obstetrics, University of Freiburg Medical Center, Freiburg, Germany*
J.S. GREENBERGER, *Department of Radiation Oncology, University of Pittsburgh, Pittsburgh, PA, USA*
J.W. GREINER, *National Cancer Institute, NIH, Bethesda, MD, USA*
C.J. GRUBBS, *Department of Nutrition Sciences, University of Alabama, Birmingham, AL, USA*
F. GUADAGNI, *Regina Elena Cancer Institute, Rome, Italy*
K.A. HAHN, *Gulf Coast Veterinary Oncology, Houston, TX, USA*
F. HALBERG, *Chronobiology Laboratories, University of Minnesota, Minneapolis, MN, USA*
T. HAMAOKA, *Department of Oncology, Osaka University, Graduate School of Medicine, Osaka, Japan*
R.R. HARDY, *Fox Chase Cancer Center, Philadelphia, PA, USA*
J. HAU, *Department of Physiology, University of Uppsala, Sweden*
G.H. HEPPNER, *Department of Immunology, Michigan Cancer Foundation, Detroit, MI, USA*
H.R. HUBBELL, *Department of Neoplastic Diseases, Hahnemann University, Philadelphia, PA, USA*
S.A. IMAM, *Huntington Medical Research Institutes, Gene Therapy Program, Pasadena, CA, USA*
C.G. IOANNIDES, *Department of Gynecologic Oncology, M.D. Anderson Cancer Center, Houston, TX, USA*
A. ITO, *Department of Oncology, Res. Inst. for Nuclear Medicine and Biology, Hiroshima University, Hiroshima, Japan*
J.R. IZBICKI, *Chirurgische Universitätsklinik, Universitätskrankenhaus Hamburg, Germany*
K.S. JEONG, *Department of Pathology, College Vet. Medicine, Kyungpook National University, Daegu, S. Korea*
H.E. KAISER, *International Society for the Study of Comparative Oncology, Baltimore, MD, USA*
M. KOUTSILIERIS, *Department of Experimental Physiology, University of Athens Medical School, Athens, Greece*

- G.R.F. KRUEGER, *Department of Anatomy II, Center for Anatomy, The University of Cologne Medical School, Cologne, Germany*
T. KUBOTA, *Department of Surgery, Keio University School of Medicine, Tokyo, Japan*
S.A. LAMPRECHT, *Strang Cancer Research Laboratory, New York, NY, USA*
G. LANDBERG, *Department of Laboratory Medicine, Lund University, Lund, Sweden*
J. LEROY, *IRCAD, University of Strasbourg, France*
B. LÉYLAND-JONES, *Department of Oncology, McGill University, Montreal, Québec, Canada*
P. MADARNAS, *Department of Pathology, University of Sherbrooke, Québec, Canada*
H. MAEDA, *Department of Microbiology, Kumamoto University, Kumamoto, Japan*
M. MAREEL, *Laboratory of Experimental Cancerology, State University, Ghent, Belgium*
J. MATTERN, *Institute of Experimental Pathology, German Cancer Research Center, Heidelberg, Germany*
D. P. MIKHAILIDIS, *Department of Chemical Pathology, The Royal Free Hospital, London, UK*
H. C. MORSE III, *Lab. of Immunopathology, National Institutes of Health, Bethesda, MD, USA*
N. MOTOHASHI, *Department of Medicinal Chemistry, Meiji Pharmaceutical University, Tokyo, Japan*
H. NAGASAWA, *Experimental Animal Research Laboratory, Meiji University, Kanagawa, Japan*
R. M. NAGLER, *Department of Oral and Maxillofacial Surgery, Rambam Medical Center, Haifa, Israel*
R. NARAYANAN, *Department of Biological Sciences, Florida Atlantic University, Boca Raton, FL, USA*
K. NILSSON, *Tumor Biology Laboratory, Department of Pathology, University Hospital, Uppsala, Sweden*
K. R. NORUM, *Institute for Nutrition Research, Universiteit i Oslo, Oslo, Norway*
R. F. NOVAK, *Institute of Chemical Toxicology, Wayne State University, Detroit, MI, USA*
J. O'KELLY, *Division of Hematology-Oncology, Cedar Sinai Medical Center, UCLA School of Medicine, Los Angeles, CA, USA*
M. PAGÉ, *Dept. de Biologie, Division de Biochimie, Université Laval, Québec, Canada*
P. PANTAZIS, *Department of Biology, Lab. of Molecular Cancer Biology, University of Miami, Coral Gables, FL, USA*
M.-F. POUPON, *Institut Curie, Section de Recherche, Paris, France*
M. PSALLIDOPOULOS, *Institute of Biomedical Research, Athens, Greece*
C.A. RUBIO, *Karolinska Institute, Pathology Research Laboratory, Stockholm, Sweden*
G.R. RUTTEMAN, *Department of Clinical Sciences of Companion Animals, University of Utrecht, Utrecht, The Netherlands*
H. SAKAGAMI, *Department of Dental Pharmacology, Meikai University School of Dentistry, Saitama, Japan*
G. SAVA, *Institutes of Biological Research, Fondazione Callario-Onlus, Trieste, Italy*
K.J. SCANLON, *Keck Graduate Institute, Claremont, CA, USA*
D. SCHIFFER, *Department of Neuroscience, University of Turin, Torino, Italy*
L.D. SHULTZ, *The Jackson Laboratory, Bar Harbor, ME, USA*
G. SICA, *Istituto di Istologia ed Embriologia, Università Cattolica del Sacro Cuore, Roma, Italy*
R.M. SNAPKA, *Department of Radiology, The Ohio State University, Columbus, OH, USA*
G.-I. SOMA, *Institutes for Health Sciences, Tokushima Bunri University, Tokushima, Japan*
T.A. SPRINGER, *Department of Pathology, Harvard Medical School, Boston, MA, USA*
K. TAKAGI, *Kennan Takagi Clinic, Kumamoto, Japan*
T. TAKAHASHI, *Aichi Cancer Center, Nagoya, Japan*
N. TANAKA, *Research Institute, Daiichi Seiyaku Co. Ltd., Tokyo, Japan*
H. TAPIERO, *Department of Cellular and Molecular Pharmacology, I.C.I.G., Villejuif, France*
K.D. TEW, *Fox Chase Cancer Center Philadelphia, PA, USA*
T. TOGE, *Research Institute for Radiation Biology, Hiroshima University, Hiroshima, Japan*
G.C. TORRE, *Department of Obstetrics and Gynecology, Albenga (SV), Italy*
B. TRIBUKAIT, *Department of Medical Radiobiology, Karolinska Hospital, Stockholm, Sweden*
C. TROPÉ, *Radiumhospitalet, Oslo, Norway*
P. TSONIS, *Department of Biology, University of Dayton, OH, USA*
T. TSURUO, *Japan Foundation Cancer Research, Cancer Chemotherapy Center, Tokyo, Japan*
J. VADGAMA, *Department of Internal Medicine, Charles E. Drew University of Medicine and Science, Los Angeles, CA, USA*
A. VECCHIONE, *Università degli Studi di Roma "La Sapienza", Rome, Italy*
N. WATANABE, *Department of Clinical Laboratory Medicine, Sapporo Medical University School of Medicine, Sapporo, Japan*
W. WEBER, *Clinical Cancer Etiology Unit, Basel, Switzerland*
L.M. WEINER, *Fox Chase Cancer Center, Philadelphia, PA, USA*
J.A. WERNER, *Department of Otolaryngology, Head and Neck Surgery, Marburg, Germany*
H. YAMAUE, *Second Department of Surgery, Wakayama Medical University, School of Medicine, Wakayama, Japan*
H. YOSHIDA, *Department of Pathology, Kagoshima University, Kagoshima, Japan*
O. YUGE, *Department of Anesthesiology, Hiroshima University School of Medicine, Hiroshima, Japan*