

## Editorial Board

- N.J. AGNANTIS, *Department of Pathology, University of Ioannina, Ioannina, Greece*  
D. ANDERSON, *Department of Biomedical Sciences, University of Bradford, Bradford, West Yorkshire, UK*  
V. BARAK, *Department of Oncology, Hadassah University Hospital, Jerusalem, Israel*  
M.H. BARCELLOS-HOFF, *Life Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA*  
Y. BECKER, *Department of Molecular Biology, Hebrew University of Jerusalem, Jerusalem, Israel*  
K. BEIER, *Department of Histology, University of Basel, Basel, Switzerland*  
S. BEN-EFRAIM, *Department of Human Microbiology, University of Tel-Aviv, Tel-Aviv, Israel*  
M. BERGQVIST, *Department of Oncology, Radiology and Clinical Immunology, University Hospital, Uppsala, Sweden*  
R. BJERKVIK, *Department of Anatomy and Cell Biology, University of Bergen, Bergen, Norway*  
B. BODEY, *Department of Pathology, University of Southern California, Reseda, CA, USA*  
D.A. BUTTERFIELD, *Department of Biological and Physical Chemistry, University of Kentucky, Lexington, KY, USA*  
M. CARAGLIA, *Department of Experimental Oncology, National Institute of Tumours Fondazione G. Pascale, Naples, Italy*  
P. CHANDRA, *Department of Molecular Biology, Frankfurt University, Frankfurt am Main, Germany*  
J.-G. CHUNG, *Department of Medicine, China Medical College, Taichung, Taiwan, ROC*  
L.A. COHEN, *Institute for Cancer Prevention, Valhalla, NY, USA*  
A.I. CONSTANTINOU, *Department of Biological Sciences, University of Cyprus, Nicosia, Cyprus*  
J.G. CORY, *Department of Biochemistry and Molecular Biology, East Carolina University, Greenville, NC, USA*  
T. DALIANIS, *Department of Pathology-Oncology, Karolinska Institute, Stockholm, Sweden*  
I. DAMJANOV, *Department of Pathology, The University of Kansas, Kansas City, KS, USA*  
G. DELICONSTANTINOS, *Department of Experimental Physiology, University of Athens Medical School, Athens, Greece*  
D.T. DENHARDT, *Department of Cell Biology and Neurosciences, Rutgers University, Piscataway, NJ, USA*  
W. DEN OTTER, *Department of Biochemistry, University of Utrecht, Utrecht, The Netherlands*  
K. DE MEIRLEIR, *Department of Human Physiology and Medicine, Faculty of Physical Education and Physiotherapy, Vrije Universiteit Brussel, Brussels, Belgium*  
L. DE RIDDER, *Department of Anatomy, Embryology and Histology, University of Ghent, Ghent, Belgium*  
E.P. DIAMANDIS, *Department of Pathology and Laboratory Medicine, University of Toronto, Toronto, Ontario, Canada*  
J.F. DORÉ, *Centre Léon Berard, Lyon, France*  
T. EFFERTH, *Helmholtz Graduate School for Cancer Research, German Cancer Research Center, Heidelberg, Germany*  
I. EMBER, *Department of Public Health, University of Pécs Medical School, Pécs, Hungary*  
W. ENGSTRÖM, *Department of Molecular Biosciences, Swedish University of Agricultural Sciences, Uppsala, Sweden*  
M. ESKELINEN, *Department of Surgery, University Hospital of Kuopio, Kuopio, Finland*  
J.A. FERNANDEZ-POL, *Metalloproteomics, LLC, Chesterfield, MO, USA*  
S. FERRONE, *Department of Immunology, Roswell Park Cancer Institute, Buffalo, NY, USA*  
G. FIORENTINI, *Department of Oncology, San Giuseppe – Antica Sede Hospital, Empoli (FI), Italy*  
P.B. FISHER, *Department of Pathology, Neurosurgery and Urology, Columbia University, New York, NY, USA*  
I. FREITAS, *Dipartimento di Biologia Animale, University of Pavia, Pavia, Italy*  
M. FRIEDRICH, *Department of Obstetrics and Gynecology, Klinikum Krefeld, Krefeld, Germany*  
R.E. FRIEDRICH, *Department of Oral and Maxillofacial Surgery, Eppendorf University Hospital, Hamburg, Germany*  
R. GANAPATHI, *Taussig Cancer Center, The Cleveland Clinic Foundation, Cleveland, OH, USA*  
D.H. GILDEN, *Department of Neurology, University of Colorado, Denver, CO, USA*  
G. GITSCH, *Department of Gynecology and Obstetrics, University of Freiburg Medical Center, Freiburg, Germany*  
J.S. GREENBERGER, *Department of Radiation Oncology, University of Pittsburgh, Pittsburgh, PA, USA*  
J.W. GREINER, *National Cancer Institute, NIH, Bethesda, MD, USA*  
C.J. GRUBBS, *Department of Nutrition Sciences, University of Alabama, Birmingham, AL, USA*  
F. GUADAGNI, *IRCCS San Raffaele, Rome, Italy*  
F. HALBERG, *Chronobiology Laboratories, University of Minnesota, Minneapolis, MN, USA*  
T. HAMAOKA, *Department of Oncology, Osaka University, Graduate School of Medicine, Osaka, Japan*  
R.R. HARDY, *Fox Chase Cancer Center, Philadelphia, PA, USA*  
J. HAU, *Department of Comparative Medicine, University of Copenhagen, Denmark*  
M. HAUER-JENSEN, *Arkansas Cancer Research Center, University of Arkansas Medical Sciences, Little Rock, AR, USA*  
G.H. HEPPNER, *Department of Immunology, Michigan Cancer Foundation, Detroit, MI, USA*  
K. HIBI, *Department of Surgery, Showa University Fujigaoka Hospital, Yokohama, Japan*  
S.A. IMAM, *Huntington Medical Research Institutes, Gene Therapy Program, Pasadena, CA, USA*  
C.G. IOANNIDES, *Department of Gynecologic Oncology, The University of Texas M.D. Anderson Cancer Center, Houston, TX, USA*  
J.R. IZBICKI, *Chirurgische Universitätsklinik, Universitätskrankenhaus Hamburg, Germany*  
K.S. JEONG, *Department of Pathology, College of Veterinary Medicine, Kyungpook National University, Daegu, S. Korea*

- M. KOUTSILIERIS, *Department of Experimental Physiology, University of Athens Medical School, Athens, Greece*  
G.R.F. KRUEGER, *Department of Anatomy II, Center for Anatomy, The University of Cologne Medical School, Cologne, Germany*  
T. KUBOTA, *Department of Surgery, Keio University School of Medicine, Tokyo, Japan*  
S.A. LAMPRECHT, *Strang Cancer Research Laboratory, New York, NY, USA*  
G. LANDBERG, *Department of Laboratory Medicine, Lund University, Lund, Sweden*  
I. LELONG-REBEL, *Lab. de Cancérologie Exp. et de Radiobiologie, IRCAD, Strasbourg, France*  
J. LEROY, *IRCAD, University of Strasbourg, France*  
B. LÉYLAND-JONES, *Winship Cancer Institute, Emory University School of Medicine, Atlanta, GA, USA*  
W. LICHTENEGGER, *Charité Campus Virchow-Klinikum, Berlin, Germany*  
P. MADARNAS, *Department of Pathology, University of Sherbrooke, Québec, Canada*  
H. MAEDA, *Faculty of Pharmaceutical Sciences, Sojo University, Kumamoto, Japan*  
M. MAREEL, *Laboratory of Experimental Cancerology, State University, Ghent, Belgium*  
G. MARTORANA, *Policlinico S. Orsola Malpighi, University of Bologna, Bologna, Italy*  
D. P. MIKHAILIDIS, *Department of Chemical Pathology, The Royal Free Hospital, London, UK*  
H. C. MORSE III, *Laboratory of Immunopathology, NIH, Bethesda, MD, USA*  
N. MOTOHASHI, *Department of Medicinal Chemistry, Meiji Pharmaceutical University, Tokyo, Japan*  
R. M. NAGLER, *Department of Oral and Maxillofacial Surgery, Rambam Medical Center, Haifa, Israel*  
R. NARAYANAN, *Department of Biological Sciences, Florida Atlantic University, Boca Raton, FL, USA*  
K. NILSSON, *Tumor Biology Laboratory, Department of Pathology, University Hospital, Uppsala, Sweden*  
K. R. NORUM, *Institute for Nutrition Research, University of Oslo, Oslo, Norway*  
R. F. NOVAK, *Institute of Chemical Toxicology, Wayne State University, Detroit, MI, USA*  
J. O'KELLY, *Division of Hematology-Oncology, Cedar Sinai Medical Center, UCLA School of Medicine, Los Angeles, CA, USA*  
Y. OKADA, *Department of Cell Physiology, National Institute for Physiological Sciences, Okazaki, Japan*  
M. PAGÉ, *Département de Biologie, Division de Biochimie, Université Laval, Québec, Canada*  
P. PANTAZIS, *Department of Biochemistry and Molecular Biology, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA*  
M.-F. POUPON, *Institut Curie, Section de Recherche, Paris, France*  
F.M. ROBERTSON, *Department of Experimental Therapeutics, The University of Texas M.D. Anderson Cancer Center, Houston, TX, USA*  
D. RUBELLO, *Department of Nuclear Medicine, PET Unit, S. Maria della Misericordia Hospital, IOV, Rovigo, Italy*  
C.A. RUBIO, *Karolinska Institute, Pathology Research Laboratory, Stockholm, Sweden*  
G.R. RUTTEMAN, *Department of Clinical Sciences of Companion Animals, University of Utrecht, Utrecht, The Netherlands*  
H. SAKAGAMI, *Department of Dental Pharmacology, Meikai University School of Dentistry, Saitama, Japan*  
G. SAVA, *Institutes of Biological Research, Fondazione Callario-Onlus, Trieste, Italy*  
K.J. SCANLON, *International BioScience, Altadena, CA, USA*  
D. SCHIFFER, *Neuro-bio-oncology Center, Policlinico di Monza Foundation, University of Turin, Vercelli, Italy*  
L.D. SHULTZ, *The Jackson Laboratory, Bar Harbor, ME, USA*  
G. SICA, *Istituto di Istologia ed Embriologia, Università Cattolica del Sacro Cuore, Roma, Italy*  
R.M. SNAPKA, *Department of Radiology, The Ohio State University, Columbus, OH, USA*  
G.-I. SOMA, *Institutes for Health Sciences, Tokushima Bunri University, Tokushima, Japan*  
T.A. SPRINGER, *Department of Pathology, Harvard Medical School, Boston, MA, USA*  
T. TAKAHASHI, *Aichi Cancer Center, Nagoya, Japan*  
N. TANAKA, *Research Institute, Daiichi Seiyaku Co. Ltd., Tokyo, Japan*  
H. TAPIERO, *Biomedicine & Pharmacotherapy, ICIG, Hopital Paul-Brousse, Villejuif, France*  
K.D. TEW, *Fox Chase Cancer Center, Philadelphia, PA, USA*  
G.C. TORRE, *Centro Ippocrate, Finale Ligure (SV), Italy*  
B. TRIBUKAIT, *Department of Medical Radiobiology, Karolinska Hospital, Stockholm, Sweden*  
C. TROPÉ, *Radiumhospitalet, Oslo, Norway*  
P. TSONIS, *Department of Biology, University of Dayton, OH, USA*  
T. TSURUO, *Japan Foundation for Cancer Research, Cancer Chemotherapy Center, Tokyo, Japan*  
J. VADGAMA, *Department of Internal Medicine, Charles E. Drew University of Medicine and Science, Los Angeles, CA, USA*  
A. VECCHIONE, *Università degli Studi di Roma "La Sapienza", Rome, Italy*  
N. WATANABE, *Department of Clinical Laboratory Medicine, Sapporo Medical University School of Medicine, Sapporo, Japan*  
W. WEBER, *Clinical Cancer Etiology Unit, Basel, Switzerland*  
L.M. WEINER, *Fox Chase Cancer Center, Philadelphia, PA, USA*  
J.A. WERNER, *Department of Otolaryngology, Head and Neck Surgery, Marburg, Germany*  
H. YAMAUE, *Second Department of Surgery, Wakayama Medical University, School of Medicine, Wakayama, Japan*  
S. YLÄ-HERTTUALA, *Department of Molecular Medicine, A.I. Virtanen Institute, University of Kuopio, Finland*  
H. YOSHIDA, *Department of Pathology, Kagoshima University, Kagoshima, Japan*