

# Inhibition of NO Production in LPS-stimulated Mouse Macrophage-like Cells by Benzo[*b*]cyclohept[*e*][1,4]oxazine and 2-Aminotropone Derivatives

AKINA SUGA<sup>1</sup>, TAICHI NARITA<sup>1</sup>, LI ZHOU<sup>2</sup>, HIROSHI SAKAGAMI<sup>3</sup>,  
KAZUE SATOH<sup>4</sup> and HIDETSUGU WAKABAYASHI<sup>1</sup>

<sup>1</sup>Faculty of Science, Josai University, Sakado, Saitama;

<sup>2</sup>Meikai Pharmaco-Medical Laboratory (MPL) and <sup>3</sup>Division of Pharmacology,

Department of Diagnostic and Therapeutic Sciences, Meikai University School of Dentistry, Saitama;

<sup>4</sup>Department of Anatomy, School of Medicine, Showa University, Tokyo, Japan

**Abstract.** The aim of this study was to investigate whether a total of twenty benzo[*b*]cyclohept[*e*][1,4]oxazines and their *S*-analogs, and 2-aminotropone derivatives affect the function of activated macrophages. These compounds inhibited the production of pro-inflammatory substances such as nitric oxide (NO) by lipopolysaccharide (LPS)-activated mouse macrophage-like RAW264.7 cells to different extents. Among them, benzo[*b*]cyclohept[*e*][1,4]oxazin-6(11*H*)-one [5] and 7-bromo-2-(4-hydroxyanilino)tropone [16] showed the highest inhibitory effects at concentrations that did not affect cellular viability (selectivity index=74.89 and 54.15, respectively). Western blot and RT-PCR analyses showed that [16] inhibited the expression of both inducible NO synthase (*iNOS*) and cyclooxygenase (COX)-2 at both protein and mRNA levels, whereas [5] inhibited only *iNOS* protein expression. Electron-spin resonance (ESR) spectroscopy revealed that both [5] and [16] scavenged nitric oxide (generated from NOC-7) and superoxide anion (generated by HX-XOD reaction) only at much higher concentration. These data suggest that [16] but not [5] exerts its anti-inflammatory action against macrophages via the inhibition of *iNOS* and COX-2 protein expressions.

Heterocyclic compounds are known to display diverse biological activities (1). Hinokitiol and its related derivatives with a tropolone skeleton (2-4) have been reported to exhibit various

biological activities such as antimicrobial (5), antifungal (6) and phytogrowth-inhibitory activity (7, 8), cytotoxic effects on mammalian tumor cells (9, 10), and inhibitory effects on catechol-*O*-methyltransferase (11) and metalloproteases (5). Hinokitiol acetate did not exhibit cytotoxic activity (10), antimicrobial activity nor metalloprotease inhibition (5), suggesting that these biological effects of hinokitiol-related compounds may result from the formation of metal chelates by the carbonyl group at C-1 with the hydroxyl group at C-2 in the tropone skeleton.

A total of 173 azulene, tropolone and azulenequinone-related compounds have been investigated for their ability to induce tumor-specific cytotoxicity and to inhibit nitric oxide (NO) and prostaglandin (PG) E<sub>2</sub> production by lipopolysaccharide (LPS)-stimulated mouse macrophage-like RAW264.7 cells (12-25). We have previously reported that tropolone, 2,4-dibromo-7-methoxytropone (14), diethyl 2-chloroazulene-1,3-dicarboxylate, 1,3-difluoroazulene (12), 3,7-dibromo-1,5-azulenequinone and 3-morpholino-1,5-azulenequinone (16) potently inhibited NO production by LPS-activated RAW264.7 cells. However, the extent of inhibition of inducible NO synthase (*iNOS*) expression differed considerably from compound to compound (12, 14, 16). Furthermore, among these 6 compounds, only diethyl 2-chloroazulene-1,3-dicarboxylate effectively inhibited PGE<sub>2</sub> production by LPS-stimulated RAW264.7 cells. These data demonstrated the lack of correlation between the extent of inhibition of *iNOS* protein expression by tropolone or azulene derivatives and that of PGE<sub>2</sub> (24).

Differing from unreactive heterocyclic-annulated tropylium compounds, benzo[*b*]cyclohept[*e*][1,4]oxazines and their *S*-analogs are generally very reactive, especially towards 1,4-difunctional nucleophiles such as *O*-phenylenediamine, ethylenediamine and their *S*- and *O*-analogs (26). We have recently reported the tumor-specific cytotoxicity of a total of

Correspondence to: Dr. Hidetsugu Wakabayashi, Faculty of Science, Josai University, Sakado, Saitama 350-0295, Japan. Tel: +81 0492717959, Fax: +81 0492717985, e-mail: hwaka@josai.ac.jp

**Key Words:** Benzo[*b*]cyclohept[*e*][1,4]oxazines, 2-aminotropones, RAW264.7 cells, macrophage, NO, COX-2, anti-inflammatory activity.




Figure 1. Chemical structure of benzo[b]cyclohept[e][1,4]benzoxazines and their S-analogs [1-8], and 2-aminotropone [9-20].

twenty benzo[b]cyclohept[e][1,4]oxazines and their S-analogs, and 2-aminotropone derivatives (25). In order to confirm the anti-inflammatory potency of these twenty compounds (Figure 1), we investigated whether they inhibit NO production by LPS-stimulated RAW264.7 cells. In the arachidonic acid (AA) cascade, cyclooxygenase (COX) catalyzes the rate-limiting reactions for prostaglandin (PG) synthesis comprising the *bis*-cyclooxygenation of AA to form PGG<sub>2</sub> and the peroxidative reduction of this intermediate to PGH<sub>2</sub> (27). Since COX has two isoforms, COX-1 (constitutive isoform) and COX-2 (inducible isoform) (28), we also investigated whether these compounds inhibit the expression of COX-2 as well as iNOS at protein and mRNA levels.

## Materials and Methods

**Materials.** The following chemicals and reagents were obtained from the indicated companies: Dulbecco's modified Eagle's medium (DMEM) (Gibco BRL, Grand Island, NY, USA); fetal bovine serum (FBS) (JRH Bioscience, Lenexa, KS, USA); LPS from *Escherichia coli* (Serotype 0111:B4), 3-(4,5-dimethylthiazol-

Table I. Benzo[b]cyclohepta[e][1,4]benzoxazines and their S-analogs [1-8], and 2-aminotropolones [9-20] synthesized in this study.

Compd.	Name
[1]	Benzo[b]cyclohepta[e][1,4]thiazin-10(11H)-one
[2]	7,9-Dibromobenzo[b]cyclohepta[e][1,4]thiazin-10(11H)-one
[3]	Benzo[b]cyclohepta[e][1,4]oxazin-10(11H)-one
[4]	7,9-Dibromobenzo[b]cyclohepta[e][1,4]oxazin-10(11H)-one
[5]	Benzo[b]cyclohepta[e][1,4]oxazin-6(11H)-one
[6]	4-Formylphenoxazine
[7]	1-Formylphenoxazine
[8]	8-Chloro-1-formylphenoxazine
[9]	2-(2-Hydroxyanilino)tropone
[10]	7-Bromo-2-(2-hydroxyanilino)tropone
[11]	5-Bromo-2-(2-hydroxyanilino)tropone
[12]	5,7-Dibromo-2-(2-hydroxyanilino)tropone
[13]	7-Bromo-2-(2-methoxyanilino)tropone
[14]	5-Bromo-2-(2-methoxyanilino)tropone
[15]	5,7-Bromo-2-(2-methoxyanilino)tropone
[16]	7-Bromo-2-(4-hydroxyanilino)tropone
[17]	7-Bromo-2-(4-methoxyanilino)tropone
[18]	5-Isopropyl-2-(2-hydroxyanilino)tropone
[19]	6-Isopropyl-2-(2-hydroxyanilino)tropone
[20]	4-Isopropyl-2-(2-hydroxyanilino)tropone

2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma Chem Co., St Louis, MO, USA); dimethyl sulfoxide (DMSO) (Wako Pure Chem. Ind., Osaka, Japan); 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (carboxy-PTIO; a spin trap agent), 1-hydroxy-2-oxo-3-(N-3-methyl-3-aminopropyl)-3-methyl-1-triazene (NOC-7; an NO generator), diethylenetriaminepentaacetic acid (DETAPAC), 5,5-dimethyl-1-pyrroline-N-oxide (DMPO) (Dojin, Kumamoto, Japan).

**Synthesis of benzo[b]cyclohepta[e][1,4]oxazines and their S-analogs, and 2-aminotropolones.** Benzo[b]cyclohepta[e][1,4]oxazines and their S-analogs [1-8] and 2-aminotropone derivatives [9-20] (Figure 1, Table I) were synthesized as described elsewhere (25).

**Cell culture.** RAW264.7 cells (29) were cultured as an adherent monolayer culture in DMEM supplemented with 10% heat-inactivated FBS under a humidified 5% CO<sub>2</sub> atmosphere, as reported elsewhere (23). When the cells were grown to confluence, they were detached by being splashed with the medium through the pipette, and seeded at lower cell density. We did not collect the cells by scraping them out from the plate with a rubber policeman, since the scraping procedure damaged the cells.

**Determination of viable cell number.** RAW264.7 cells were inoculated at 0.3×10<sup>6</sup>/ml (100 µl) in a 96-microwell plate (Becton Dickinson Labware, NJ, USA) and incubated for 24 hours. Near confluent cells were treated for a further 24 hours with the indicated concentrations of test compound in phenol red-free DMEM supplemented with 10% FBS in the presence or absence of LPS (0.1 µg/ml). The culture supernatant was collected for the determination of NO as described below. The attached cells were used for the determination of viable cell number by MTT method. In brief, cells were incubated for 60 minutes with 0.2 mg/ml of MTT. After removal of the medium, the

Table II. Effect of benzo[b]cyclohepta[e][1,4]oxazine and 2-aminotropone derivatives on NO production by LPS-stimulated RAW264.7 cells.


Compd.	CC <sub>50</sub> (µM)	IC <sub>50</sub> (µM)	SI
[1]	>100.00±0.00	14.7±10.07	6.80
[2]	>100.00±0.00	5.47±3.74	18.28
[3]	>125.00±0.00	14.2±10.80	8.80
[4]	>12.5.00±0.00	3.16±2.18	3.96
[5]	342.77±88.66	4.58±5.23	74.89
[6]	>125.00±0.00	21.66±17.99	5.77
[7]	>50.00±0.00	13.36±9.67	3.74
[8]	>100.00±0.00	11.17±9.30	8.95
[9]	500.00±0.00	37.36±27.09	13.38
[10]	116.19±11.01	11.06±7.67	10.51
[11]	142.13±57.46	13.76±7.96	10.33
[12]	84.03±32.72	6.57±5.67	12.79
[13]	155.76±29.05	12.33±7.22	12.63
[14]	>125.00±0.00	28.94±22.96	4.32
[15]	>100.00±0.00	13.48±6.26	7.42
[16]	128.42±18.72	2.37±1.37	54.15
[17]	139.84±18.94	11.27±5.09	12.40
[18]	112.24±11.06	9.25±2.78	12.13
[19]	123.45±35.14	11.03±10.35	11.19
[20]	134.48±35.69	15.02±10.73	8.96

CC<sub>50</sub>: 50% Cytotoxic concentration; IC<sub>50</sub>: concentration at which LPS-stimulated NO production was inhibited by 50%; SI: selectivity index (CC<sub>50</sub>/IC<sub>50</sub>). Each value represents the mean±S.D. from at least 3 independent experiments.

cells were lysed with DMSO, and the absorbance at 540 nm was recorded using a microplate reader (23).

**Assay for NO production.** RAW264.7 cells were treated with test compounds as described above, and the concentration of NO released from the cells was determined by Greiss reagent, using the standard curve of NO<sub>2</sub><sup>-</sup>. To eliminate the interaction between sample and Greiss reagent, we also measured the NO concentration in the culture medium without the cells and subtracted this value from that obtained with the cells. The concentration that inhibited the LPS-stimulated NO production by 50% (50% inhibitory concentration: IC<sub>50</sub>) was determined from the dose-response curve (23). The efficacy of inhibition of NO production was evaluated by the selectivity index (SI), which was calculated using the following equation: SI=CC<sub>50</sub>/IC<sub>50</sub>, where CC<sub>50</sub> was the cytotoxic concentration causing death of 50% of the population of cells.

**Assay for iNOS protein expression.** RAW264.7 cells were inoculated at 1.0×10<sup>6</sup>/ml (100 µl) in 96-well plates (Becton Dickinson) and incubated for 24 hours. Near-adherent cells were treated for 24 hours with different concentrations of test sample. The concentrations were selected by CC<sub>50</sub> in the presence or absence of 0.1 µg/ml LPS. The cell pellets were lysed with 50 µl of lysis buffer [10 mM Tris-HCl (pH 7.6), 1% Triton® X-100, 150 mM NaCl, 5 mM EDTA-2Na and 2 mM PMSF] for 10 minutes on ice. The cell lysates were centrifuged at 16,000×g for 20 minutes at 4°C to remove insoluble materials and the supernatant was collected. The protein concentrations of the supernatant were measured by Protein Assay Kit (Bio Rad, Hercules, CA, USA). Equal amounts of the protein



**Figure 2.** Effect of benzo[*b*]cyclohept[e][1,4]oxazin-6(11H)-one [5] and 7-bromo-2-(4-hydroxyanilino)tropone [16] on the LPS-stimulated expression of iNOS and COX-2 proteins. RAW264.7 cells ( $100 \times 10^4$  cells/ml, 1 ml) were inoculated on 24-well plate and incubated for 24 hours. RAW264.7 cells were then treated for 24 hours without (control) or with  $\times 1/2$ ,  $\times 1$ ,  $\times 2$  or  $\times 4$  of  $IC_{50}$  (concentration that inhibited the LPS-stimulated NO production by 50%) of benzo[*b*]cyclohept[e][1,4]oxazin-6(11H)-one [5] or 7-bromo-2-(4-hydroxyanilino)tropone [16] in the presence or absence of LPS (0.1  $\mu$ g/ml) in the presence or absence of LPS (0.1  $\mu$ g/ml), and then the expression of iNOS and COX-2 proteins were monitored by Western blot analysis.

from cell lysates (10  $\mu$ g) were mixed with 2×sodium dodecyl sulfate (SDS)-sample buffer [0.1 M Tris-HCl (pH 6.8), 20% glycerol, 4% SDS, 0.01% bromophenol blue, 1.2% 2-mercaptoethanol], boiled for 10 minutes, and applied to SDS-8% polyacrylamide gel electrophoresis, then transferred for 90 minutes to polyvinylidene difluoride (PVDF) membrane. The membranes were blocked with 5% skim milk in phosphate-buffered saline [PBS (-)] plus 0.05% Tween 20 for 60 minutes and incubated for 90 minutes at room temperature with anti-iNOS or anti-COX-2 (dilution: 1:1,000 and 1:4,000, respectively, Santa Cruz Biotechnology, Delaware, CA, USA) or anti-actin antibody (1:6,000; Sigma Chem Co.), and then incubated with horseradish peroxidase-conjugated anti-goat, anti-rabbit or anti-mouse IgG for 60 minutes at room temperature. Immunoblots were developed with Western Lighting™ Chemiluminescence Reagent Plus (21).

**Assay for iNOS and COX-2 mRNA expression.** RAW264.7 cells were inoculated at  $1.0 \times 10^6$ /ml (100  $\mu$ l) in 96-well plates and incubated for 24 hours. Near-confluent cells were treated for 24 hours with different concentrations of test compound in the presence or absence of LPS (0.1  $\mu$ g/ml). The expression levels of iNOS and COX-2 mRNA were measured by RT-PCR. The primer sequence and size was: (i) iNOS: PCR product size 496 bp, sense primer 5'-CCCTTCCGAAGTTCTGGCAGCAGC-3', antisense primer 5'-GGCTGTCAGAGCCTCGTGGCTTG-3'; (ii) COX-2: Product size 371 bp, sense primer 5'-TTTGTTGAGTCATTACCAAGACAGAT-3', antisense primer 5'-CAGTATTGAGGAGAACAGA TGGGATT-3'; (iii) G3PDH: Product size 452 bp, sense primer 5'-ACCACAGTCCATGCCATCAC-3', antisense primer 5'-TCCACC ACCCTGTTGCTGTA-3' (21). After amplification, each reaction mixture was analyzed by 2% agarose gel electrophoresis and visualized by ethidium bromide staining under UV light. The relative amount of PCR product was normalized to the corresponding amount of G3PDH PCR product.

**Radical-scavenging activity.** The radical intensity of test samples was determined at 25°C, using electron-spin resonance (ESR) spectroscopy (JEOL JES REIX Tokyo, Japan; X-band, 100 kHz modulation frequency) (21, 23). The radical intensity of NO produced from the reaction mixture of 20  $\mu$ M carboxy-PTIO and 60  $\mu$ M NOC-7 was determined in 0.1 M phosphate buffer (PB; pH 7.4)

in the presence of 30% DMSO (microwave power and gain were 5 mW and 400, respectively). When NOC-7 and carboxy-PTIO were mixed, NO was oxidized to  $NO_2$  and carboxy-PTIO was reduced to carboxy-PTI, which produces seven-line signals. Samples were added 3 minutes after mixing. The NO radical intensity was defined as the ratio of the signal intensity of the first peak of carboxy-PTI to that of MnO.


The superoxide anion (in the form of DMPO-OOH), produced by the HX-XOD reaction (total volume: 200  $\mu$ l) [2 mM HX in 0.1 M (pH 7.4) (PB) 50  $\mu$ l, 0.5 mM DETAPAC 20  $\mu$ l, 8% DMPO 30  $\mu$ l, sample 40  $\mu$ l, H<sub>2</sub>O 30  $\mu$ l, XOD (0.5 U/ml in PB) 30  $\mu$ l] was determined (microwave power: 16 mW, gain: 630) (30).

## Results

**Inhibition of NO production.** Twenty benzo[*b*]cyclohept[e][1,4]oxazines and their *S*-analogs, and 2-aminotropone derivatives inhibited the production of NO by LPS-activated mouse macrophage-like RAW264.7 cells to different extents. Among them, benzo[*b*]cyclohept[e][1,4]oxazin-6(11H)-one [5] and 7-bromo-2-(4-hydroxyanilino)tropone [16] showed the highest inhibitory effects at concentrations that did not affect the cellular viability (SI=74.89 and 54.15, respectively, Table II).

**Effect on iNOS and COX-2 expression.** Western blot analyses showed that both compound [5] and [16] inhibited expression of iNOS protein (Figure 2). Compound [16] inhibited COX-2 protein expression at a concentration twice and four times that of the  $IC_{50}$  (50% inhibitory concentration of LPS-stimulated NO production). On the other hand, [5] did not completely inhibit the COX-2 protein expression even at concentrations 4 times that of the  $IC_{50}$ .

RT-PCR analysis showed that [16] but not [5] inhibited iNOS and COX-2 mRNA expression at a concentration twice and four times that of the  $IC_{50}$  (Figure 3).



**Figure 3.** Effect of benzo[*b*]cyclohept[*e*][1,4]oxazin-6(11*H*)-one [5] and 7-bromo-2-(4-hydroxyanilino)tropone [16] on the LPS-stimulated expression of iNOS and COX-2 mRNA. RAW264.7 cells ( $100 \times 10^4$  cells/ml, 1 ml) were inoculated on 24-well plate and incubated for 24 hours. RAW264.7 cells were then treated for 24 hours without (control) or with  $\times 1/2$ ,  $\times 1$ ,  $\times 2$  or  $\times 4$  of IC<sub>50</sub> (concentration that inhibited the LPS-stimulated NO production by 50%) of benzo[*b*]cyclohept[*e*][1,4]oxazin-6(11*H*)-one [5] or 7-bromo-2-(4-hydroxyanilino)tropone [16] in the presence or absence of LPS (0.1  $\mu$ g/ml) in the presence or absence of LPS (0.1  $\mu$ g/ml), and then the expression of iNOS and COX-2 mRNA was monitored by RT-PCR analysis.

*NO-scavenging activity.* There was a possibility that the reduced concentration of NO in the culture medium of the cells treated with either [5] or [16] may have been due to the NO-scavenging activity of these compounds. ESR spectroscopy revealed that both [5] and [16] scavenged NO generated from NOC-7 only at much higher concentrations (30-200 times higher than the concentration of IC<sub>50</sub> for the inhibition of NO production; Figure 4A), demonstrating that this possibility was small. Since NO and O<sub>2</sub><sup>-</sup> interact with each other, we investigated whether [5] and [16] scavenge the O<sub>2</sub><sup>-</sup> generated by the HX-XOD reaction. ESR spectroscopy revealed that these compounds also scavenged O<sub>2</sub><sup>-</sup> very poorly (Figure 4B).

## Discussion

The present study demonstrated that all twenty benzo[*b*]cyclohept[*e*][1,4]oxazine and 2-aminotropone derivatives investigated showed an inhibitory effect on the NO production by LPS-activated RAW264.7 cells to different extents, and that among them, benzo[*b*]cyclohept[*e*][1,4]oxazin-6(11*H*)-one [5] and 7-bromo-2-(4-hydroxyanilino)tropone [16] were the most potent inhibitors. However, these two compounds seem to inhibit LPS-stimulated NO production by different mechanisms.

Compound [16] inhibited iNOS protein expression more efficiently than that of NO production, and therefore the inhibition of NO production by [16] was due to the inhibition

of iNOS production (Figure 2) resulting from impaired mRNA expression, not due to its NO-scavenging activity (Figure 4). Compound [16] also inhibited the expression of COX-2 protein (Figure 2) and mRNA expression (Figure 3), but at slightly higher concentrations ( $\times 2$  and  $\times 4$  of IC<sub>50</sub> of NO production). These data suggest the possible anti-inflammatory activity of [16]. Higher tumor specificity of this compound (25) makes the further investigation of [16] promising.

Similarly, the inhibition of LPS-stimulated NO production by compound [5] was due to the inhibition of the induction of iNOS protein expression, not due to its NO-scavenging activity. However, [5] failed to inhibit iNOS mRNA expression. It remains to be investigated whether [5] induces post-transcriptional modification of iNOS, and whether it induces the S-nitrosylation of COX-2 that is responsible for the decline of this enzyme's activity (31).

In conclusion, the present study suggests that the inhibition of NO production by macrophages is not an appropriate parameter for evaluating anti-inflammatory activity, supporting our recent report (24). Other markers, such as PGE<sub>2</sub> production or COX-2 expression, should be monitored in addition to NO production.

## Acknowledgements

This study was supported in part by a Grant-in-Aid from the Ministry of Education, Science, Sports and Culture of Japan (Sakagami, No. 19592156).




Figure 4. Low scavenging activity of benzo[b]cyclohept[e][1,4]oxazin-6(11H)-one [5] and 7-bromo-2-(4-hydroxyanilino)tropone [16] against NO and  $\text{O}_2^-$ . Each value represents the mean $\pm$ S.D. of triplicate assays.

## References

- 1 Heterocyclic Chemistry. Bioactive Heterocycles VI. Flavonoids and Anthocyanins in Plants and Latest Bioactive N-Heterocycles. Noboru Motohashi (ed.). Springer, 2008.
- 2 Nozoe T, Takase K, Matsumura H, Asao T, Kikuchi K and Ito S: Non-benzenoid aromatic compounds. In: Dai-Yuki Kagaku. Kotake M (ed.), Tokyo, Asakura-Shoten Publ. Co., Vol. 13, pp. 1-656, 1960 (in Japanese).
- 3 Nozoe T, Breslow R, Hafner K, Ito S and Murata I: Topics in non-benzenoid aromatic chemistry. Tokyo, Hirokawa Publ. Co., Vol. I, pp. 1-295, 1973.
- 4 Nozoe T, Breslow R, Hafner K, Ito S and Murata I: Topics in non-benzenoid aromatic chemistry. Tokyo, Hirokawa Publ. Co., Vol. II, pp 1-379, 1977.
- 5 Inamori Y, Shinohara S, Tsujibo H, Okabe T, Morita Y, Sakagami Y, Kumeda Y and Ishida N: Antimicrobial activity and metalloprotease inhibition of Hinokitiol-related compounds, the constituents of *Thujopsis dolabrata* S. and *Z. hondai* Mak. Biol Pharm Bull 22: 990-993, 1999.
- 6 Inamori Y, Sakagami Y, Morita Y, Shibata M, Sugiura M, Kumeda Y, Okabe T, Tsujibo H and Ishida N: Antifungal activity of hinokitiol-related compounds on wood-rotting fungi and their insecticidal activities. Biol Pharm Bull 23: 995-997, 2000.
- 7 Inamori Y, Nishiguchi K, Matsuo N, Tsujibo H, Baba K and Ishida N: Phytogrowth-inhibitory activities of tropolone and hinokitiol. Chem Pharm Bull 39: 2378-2381, 1991.
- 8 Sakagami Y, Inamori Y, Isoyama N, Tsujibo H, Okabe T, Morita Y and Ishida N: Phytogrowth-inhibitory activities of  $\beta$ -dolabrin and  $\gamma$ -thujaplicin, hinokitiol-related compounds and constituents of *Thujopsis dolabrata* Sieb. and Zucc. var *hondai* Makino. Biol Pharm Bull 23: 645-648, 2000.
- 9 Inamori Y, Tsujibo H, Ohnishi H, Ishii F, Mizugaki M, Aso H and Ishida N: Cytotoxic effect of hinokitiol and tropolone on the growth of mammalian cells and on blastogenesis of mouse splenic T-cells. Biol Pharm Bull 16: 521-523, 1993.

- 10 Matsumura E, Morita Y, Date T, Tsujibo H, Yasuda M, Okabe T, Ishida N and Inamori Y: Cytotoxicity of the hinokitiol-related compounds,  $\gamma$ -thujaplicin and  $\beta$ -dolabrin. *Biol Pharm Bull* 24: 299-302, 2001.
- 11 Borchardt RT: Catechol *O*-methyltransferase. 1. Kinetics of tropolone inhibition. *J Medicinal Chem* 16: 377-382, 1973.
- 12 Hashiba K, Yokoyama K, Wakabayashi H, Hashimoto K, Satoh K, Kurihara T, Motohashi N and Sakagami H: Inhibition of LPS-stimulated NO production in mouse macrophage-like cells by azulenes. *Anticancer Res* 24: 3939-3944, 2004.
- 13 Wakabayashi H, Hashiba K, Yokoyama K, Hashimoto K, Kikuchi H, Nishikawa H, Kurihara T, Satoh K, Shiota S, Saito S, Kusano S, Nakashima H, Motohashi N and Sakagami H: Cytotoxic activity of azulenes against human oral tumor cell lines. *Anticancer Res* 23: 4747-4756, 2003.
- 14 Yokoyama K, Hashiba K, Wakabayashi H, Hashimoto K, Satoh K, Kurihara T, Motohashi N and Sakagami H: Inhibition of LPS-stimulated NO production in mouse macrophage-like cells by tropolones. *Anticancer Res* 24: 3917-3922, 2004.
- 15 Wakabayashi H, Yokoyama K, Hashiba K, Hashimoto K, Kikuchi H, Nishikawa H, Kurihara T, Satoh K, Shiota S, Saito S, Kusano S, Nakashima H, Motohashi N and Sakagami H: Cytotoxic activity of tropolones against human oral tumor cell lines. *Anticancer Res* 23: 4757-4764, 2003.
- 16 Nishishiro M, Arikawa S, Wakabayashi H, Hashimoto K, Satoh K, Yokoyama K, Unten S, Kakuta H, Kurihara T, Motohashi N and Sakagami H: Inhibition of LPS-stimulated NO production in mouse macrophage-like cells by azulenequinones. *Anticancer Res* 25: 4157-4164, 2005.
- 17 Wakabayashi H, Nishishiro M, Arikawa S, Hashimoto K, Kikuchi H, Nishikawa H, Kurihara T, Terakubo S, Shoji Y, Nakashima H, Motohashi N and Sakagami H: Cytotoxic activity of azulenequinones against human oral tumor cell lines. *Anticancer Res* 25: 305-312, 2005.
- 18 Ohshima N, Akatsu Y, Yamagishi Y, Nishishiro M, Wakabayashi H, Kurihara T, Satoh K, Motohashi N and Sakagami H: Inhibition of NO production in LPS-stimulated mouse macrophage-like cells by trihaloacetylazulenes. *Anticancer Res* 26: 2921-2928, 2006.
- 19 Akatsu Y, Ohshima N, Yamagishi Y, Nishishiro M, Wakabayashi H, Kurihara T, Kikuchi H, Katayama T, Motohashi N, Shoji Y, Nakashima H and Sakagami H: Apoptosis-inducing activity of trihaloacetylazulenes against human oral tumor cell lines. *Anticancer Res* 26: 1917-1924, 2006.
- 20 Sekine T, Takahashi J, Nishishiro M, Arai A, Wakabayashi H, Kurihara T, Kobayashi M, Hashimoto K, Kikuchi H, Katayama T, Kanda Y, Kunii S, Motohashi N and Sakagami H: Tumor-specificity and type of cell death induced by trihaloacetylazulenes in human tumor cell lines. *Anticancer Res* 27: 133-144, 2007.
- 21 Takahashi J, Sekine T, Nishishiro M, Arai A, Wakabayashi H, Kurihara T, Hashimoto K, Satoh K, Motohashi N and Sakagami H: Inhibition of NO production in LPS-stimulated mouse macrophage-like cells by trihaloacetylazulene derivatives. *Anticancer Res* 28: 171-178, 2008.
- 22 Murayama H, Miyahara K, Wakabayashi H, Kurihara T, Hashimoto K, Amano O, Kikuchi H, Nakamura Y, Kanda Y, Kunii S, Motohashi N and Sakagami H: Tumor-specific cytotoxicity and type of cell death induced by benzocycloheptoxazines in human tumor cell lines. *Anticancer Res* 28: 1069-1078, 2008.
- 23 Miyahara K, Murayama H, Wakabayashi H, Kurihara T, Hashimoto K, Satoh K, Motohashi N and Sakagami H: Inhibition of LPS-stimulated NO production in mouse macrophage-like cells by benzocycloheptoxazines. *Anticancer Res* 28: 2657-2662, 2008.
- 24 Nishishiro M, Kurihara T, Wakabayashi H and Sakagami H: Effect of tropolone, azulene and azulenequinone derivatives on prostaglandin E<sub>2</sub> production by activated macrophage-like cells. *Anticancer Res* 29: 379-384, 2009.
- 25 Narita T, Suga A, Kobayashi M, Hashimoto K, Sakagami H, Motohashi N, Kurihara T and Wakabayashi H: Tumor-specific cytotoxicity and type of cell death induced by benzo[*b*]cyclohept[*e*][1,4]oxazine and 2-aminotropone derivatives. *Anticancer Res* 29: 1123-1130, 2009.
- 26 Nozoe T: Cyclohepta[*b*][1,4]benzoxazine and its related compounds. Some novel aspects in heterocyclic chemistry. *Heterocycles* 30: 1263-1306, 1990.
- 27 Odau S, Gabler C, Holder C and Einspanier R: Differential expression of cyclooxygenase 1 and cyclooxygenase 2 in the bovine oviduct. *J Endocrinol* 191: 263-274, 2006.
- 28 O'Banion MK, Winn VD and Young DA: cDNA cloning and functional activity of a glucocorticoid-regulated inflammatory cyclooxygenase. *Proc Natl Acad Sci USA* 89: 4888-4892, 1992.
- 29 Ralph R and Nakoinz: Antibody-dependent killing of erythrocyte and tumor targets by macrophage-related cell lines: enhancement by PPD and LPS. *J Immunol* 119: 950-954, 1977.
- 30 Sakagami H, Amano S, Kikuchi H, Nakamura Y, Kuroshita R, Watanabe S, Satoh K, Hasegawa H, Nomura A, Kanamoto T, Terakubo S, Nakashima H, Taniguchi S and Ohizumi T: Antiviral, antibacterial and vitamin C-synergized radical scavenging activity of *Sasa senanensis* Rehder extract. *In Vivo* 22: 471-496, 2008.
- 31 Kim SF, Huri DA and Snyder SH: Inducible nitric oxide synthase binds, *S*-nitrosylates, and activates cyclooxygenase-2. *Science* 310: 1966-1970, 2005.

*Received March 17, 2009**Revised June 3, 2009**Accepted July 2, 2009*