

## Editorial Board

- N.J. AGNANTIS, *Department of Pathology, University of Ioannina, Ioannina, Greece*  
D. ANDERSON, *Department of Biomedical Sciences, University of Bradford, Bradford, West Yorkshire, UK*  
J.P.A. BAAK, *Department of Pathology, Stavanger University Hospital, Stavanger, Norway*  
V. BARAK, *Department of Oncology, Hadassah University Hospital, Jerusalem, Israel*  
M.H. BARCELLOS-HOFF, *Department of Radiation Oncology, New York University School of Medicine, New York, NY, USA*  
Y. BECKER, *Department of Biochemistry, Faculty of Medicine, Hebrew University of Jerusalem, Jerusalem, Israel*  
K. BEIER, *Department of Histology, University of Basel, Basel, Switzerland*  
S. BEN-EFRAIM, *Department of Human Microbiology, University of Tel-Aviv, Tel-Aviv, Israel*  
M. BERGQVIST, *Department of Oncology, Radiology and Clinical Immunology, University Hospital, Uppsala, Sweden*  
R. BJERKVIG, *Norlux Neuro-Oncology, Department of Biomedicine, University of Bergen, Bergen, Norway*  
B. BODEY, *Department of Pathology, University of Southern California, Reseda, CA, USA*  
D.A. BUTTERFIELD, *Department of Biological and Physical Chemistry, University of Kentucky, Lexington, KY, USA*  
M. CARAGLIA, *Department of Experimental Oncology, National Institute of Tumours Fondazione G. Pascale, Naples, Italy*  
P. CHANDRA, *Department of Molecular Biology, Frankfurt University, Frankfurt am Main, Germany*  
J.-G. CHUNG, *Department of Medicine, China Medical College, Taichung, Taiwan, ROC*  
L.A. COHEN, *Northampton, MA, USA*  
A.I. CONSTANTINOU, *Department of Biological Sciences, University of Cyprus, Nicosia, Cyprus*  
T. DALIANIS, *Department of Pathology-Oncology, Karolinska Institute, Stockholm, Sweden*  
G. DELICONSTANTINOS, *Department of Experimental Physiology, University of Athens Medical School, Athens, Greece*  
D.T. DENHARDT, *Division of Life Sciences, Rutgers University, Bridgewater, NJ, USA*  
W. DEN OTTER, *VUMC - Department of Urology, Amsterdam, The Netherlands*  
K. DE MEIRLEIR, *Department of Human Physiology and Medicine, Faculty of Physical Education and Physiotherapy, Vrije Universiteit Brussel, Brussels, Belgium*  
L. DE RIDDER, *Department of Anatomy, Embryology and Histology, University of Ghent, Ghent, Belgium*  
E.P. DIAMANDIS, *Department of Pathology and Laboratory Medicine, University of Toronto, Toronto, Ontario, Canada*  
T. EFFERTH, *Department of Pharmaceutical Biology, Institute of Pharmacy and Biochemistry, University of Mainz, Germany*  
I. EMBER, *Department of Public Health, University of Pécs Medical School, Pécs, Hungary*  
W. ENGSTRÖM, *Department of Molecular Biosciences, Swedish University of Agricultural Sciences, Uppsala, Sweden*  
M. ESKELINEN, *Department of Surgery, University Hospital of Kuopio, Kuopio, Finland*  
J.A. FERNANDEZ-POL, *Metalloproteomics, LLC, Chesterfield, MO, USA*  
S. FERRONE, *Department of Immunology, University of Pittsburgh, Pittsburgh, PA, USA*  
G. FIORENTINI, *UOC Oncologia, Azienda Ospedaliera Marche Nord, Pesaro, Italy*  
P.B. FISHER, *Virginia Commonwealth University, School of Medicine, Richmond, VA, USA*  
I. FREITAS, *Dipartimento di Biologia Animale, University of Pavia, Pavia, Italy*  
M. FRIEDRICH, *Department of Obstetrics and Gynecology, Klinikum Krefeld, Krefeld, Germany*  
R.E. FRIEDRICH, *Department of Oral and Maxillofacial Surgery, Eppendorf University Hospital, Hamburg, Germany*  
R. GANAPATHI, *Levine Cancer Institute, Carolinas HealthCare System, Charlotte, NC, USA*  
Z. GATALICA, *Department of Pathology and Creighton Medical Laboratories, Creighton University Medical Center, Omaha, NE, USA*  
D.H. GILDEN, *Department of Neurology, University of Colorado Denver School of Medicine, Aurora, CO, USA*  
G. GITSCH, *Department of Gynecology and Obstetrics, University of Freiburg Medical Center, Freiburg, Germany*  
J.S. GREENBERGER, *Department of Radiation Oncology, University of Pittsburgh, Pittsburgh, PA, USA*

*continued*

- J.W. GREINER**, *National Cancer Institute, NIH, Bethesda, MD, USA*
- D.S. GRIDLEY**, *Department of Radiation Medicine, Radiation Research Laboratories, Loma Linda University and Medical Center, Loma Linda, CA, USA*
- C.J. GRUBBS**, *Department of Nutrition Sciences, University of Alabama, Birmingham, AL, USA*
- F. GUADAGNI**, *IRCCS San Raffaele, Rome, Italy*
- F. HALBERG**, *Chronobiology Laboratories, University of Minnesota, Minneapolis, MN, USA*
- R.R. HARDY**, *Fox Chase Cancer Center, Philadelphia, PA, USA*
- J. HAU**, *Department of Comparative Medicine, University of Copenhagen, Denmark*
- M. HAUER-JENSEN**, *Arkansas Cancer Research Center, University of Arkansas Medical Sciences, Little Rock, AR, USA*
- G.H. HEPPNER**, *Department of Immunology, Michigan Cancer Foundation, Detroit, MI, USA*
- K. HIBI**, *Department of Surgery, Showa University Fujigaoka Hospital, Yokohama, Japan*
- S.A. IMAM**, *Huntington Medical Research Institutes, Gene Therapy Program, Pasadena, CA, USA*
- C.G. IOANNIDES**, *Department of Gynecologic Oncology, The University of Texas M.D. Anderson Cancer Center, Houston, TX, USA*
- J.R. IZBICKI**, *Chirurgische Universitätsklinik, Universitätskrankenhaus Hamburg, Germany*
- K.S. JEONG**, *Department of Pathology, College of Veterinary Medicine, Kyungpook National University, Daegu, S. Korea*
- I. KISS**, *Institute of Preventive Medicine, Medical School, University of Pécs, Pécs, Hungary*
- M. KOUTSILIERIS**, *Department of Experimental Physiology, University of Athens Medical School, Athens, Greece*
- G.R.F. KRUEGER**, *Department of Anatomy II, Center for Anatomy, The University of Cologne Medical School, Cologne, Germany*
- B. KRUSLIN**, *Department of Pathology, Ljudevit Jurak University, Sestre milosrdnice University Hospital, Zagreb, Croatia*
- S.A. LAMPRECHT**, *Division of Clinical Biochemistry, Faculty of Health Sciences, Ben Gurion University of the Negev, Beer-Sheva, Israel*
- G. LANDBERG**, *Department of Laboratory Medicine, Lund University, Lund, Sweden*
- I. LELONG-REBEL**, *Lab. de Biophotonique et Pharmacologie, Université de Strasbourg, Illkirch, France*
- J. LEROY**, *IRCAD, University of Strasbourg, France*
- B. LÉYLAND-JONES**, *Edith Sanford Breast Cancer Research, Sanford Health, Sioux Falls, SD, USA*
- W. LICHTENEGGER**, *Charité Campus Virchow-Klinikum, Berlin, Germany*
- P. MADARNAS**, *Department of Pathology, University of Sherbrooke, Québec, Canada*
- H. MAEDA**, *Faculty of Pharmaceutical Sciences, Sojo University, Kumamoto, Japan*
- M. MAREEL**, *Laboratory of Experimental Cancerology, State University, Ghent, Belgium*
- G. MARTORANA**, *Policlinico S. Orsola Malpighi, University of Bologna, Bologna, Italy*
- D.P. MIKHAILIDIS**, *Department of Chemical Pathology, The Royal Free Hospital, London, UK*
- J. MOLNÁR**, *Department of Medical Microbiology and Immunobiology, Faculty of Medicine, University of Szeged, Szeged, Hungary*
- H.C. MORSE III**, *Laboratory of Immunopathology, NIH, Rockville, MD, USA*
- N. MOTOHASHI**, *Department of Medicinal Chemistry, Meiji Pharmaceutical University, Tokyo, Japan*
- R.M. NAGLER**, *Department of Oral and Maxillofacial Surgery, Rambam Medical Center, Haifa, Israel*
- R. NARAYANAN**, *Department of Biological Sciences, Florida Atlantic University, Boca Raton, FL, USA*
- K. NILSSON**, *Tumor Biology Laboratory, Department of Pathology, University Hospital, Uppsala, Sweden*
- K.R. NORUM**, *Institute for Nutrition Research, University of Oslo, Oslo, Norway*
- R.F. NOVAK**, *Shriners Hospitals for Children International, Tampa, FL, USA*
- K. OGAWA**, *Department of Surgery, Tokyo Women's Medical University Medical Center East, Tokyo, Japan*
- M. PAGÉ**, *Département de Biologie, Division de Biochimie, Université Laval, Québec, Canada*

- P. PANTAZIS**, *Department of Biochemistry and Molecular Biology, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA*
- M.-F. POUPON**, *Institut Curie, Section de Recherche, Paris, France*
- F.M. ROBERTSON**, *Department of Experimental Therapeutics, The University of Texas M.D. Anderson Cancer Center, Houston, TX, USA*
- D. RUBELLO**, *Department of Nuclear Medicine, PET Unit, S. Maria della Misericordia Hospital, IOV, Rovigo, Italy*
- C.A. RUBIO**, *Karolinska Institute, Pathology Research Laboratory, Stockholm, Sweden*
- G.R. RUTTEMAN**, *Department of Clinical Sciences of Companion Animals, University of Utrecht, Utrecht, The Netherlands*
- H. SAKAGAMI**, *Department of Dental Pharmacology, Meikai University School of Dentistry, Saitama, Japan*
- G. SAVA**, *Institutes of Biological Research, Fondazione Callerio-Onlus, Trieste, Italy*
- D. SCHIFFER**, *Neuro-bio-oncology Center, Policlinico di Monza Foundation, University of Turin, Vercelli, Italy*
- L.D. SHULTZ**, *The Jackson Laboratory, Bar Harbor, ME, USA*
- G. SICA**, *Istituto di Istologia ed Embriologia, Universita Cattolica del Sacro Cuore, Roma, Italy*
- R.M. SNAPKA**, *Department of Radiology, The Ohio State University, Columbus, OH, USA*
- G.-I. SOMA**, *Institutes for Health Sciences, Tokushima Bunri University, Tokushima, Japan*
- T.A. SPRINGER**, *Department of Pathology, Harvard Medical School, Boston, MA, USA*
- T. TAKAHASHI**, *Aichi Cancer Center, Nagoya, Japan*
- N. TANAKA**, *Department of Animal Pharmaceutical Science, Chiba Institute of Science, Chiba, Japan*
- K.D. TEW**, *Department of Pharmacology, Medical University of South Carolina, Charleston, SC, USA*
- G.C. TORRE**, *Centro Ippocrate, Vico Cosmelli 4, Finale Ligure (SV), Italy*
- B. TRIBUKAIT**, *Department of Medical Radiobiology, Karolinska Hospital, Stockholm, Sweden*
- J. VADGAMA**, *Department of Internal Medicine, Charles E. Drew University of Medicine and Science, Los Angeles, CA, USA*
- J.K. VISHWANATHA**, *Graduate School of Biomedical Sciences, University of North Texas Health Science Center, Fort Worth, TX, USA*
- N. WATANABE**, *Department of Clinical Laboratory Medicine, Sapporo Medical University School of Medicine, Sapporo, Japan*
- W. WEBER**, *Clinical Cancer Etiology Unit, Basel, Switzerland*
- L.M. WEINER**, *Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, DC, USA*
- J.A. WERNER**, *Department of Otolaryngology, Head and Neck Surgery, Marburg, Germany*
- S. YLÄ-HERTTUALA**, *Department of Molecular Medicine, A.I. Virtanen Institute, University of Kuopio, Finland*
- H. YOSHIDA**, *Department of Pathology, Kagoshima University, Kagoshima, Japan*