

Amelioration of Mucositis in Proton Therapy of Fanconi Anemia *Fanca*^{-/-} Mice by JP4-039

THOMAS J. QUINN¹, XUANFENG DING¹, XIAOQIANG LI¹, GEORGE D. WILSON¹,
KATIE BUELOW¹, ARANEE SIVANANTHAN², STEPHANIE THERMOZIER²,
ANDREW HENDERSON², MICHAEL W. EPPERLY², DARCY FRANICOLA²,
PETER WIPF³, JOEL S. GREENBERGER², CRAIG W. STEVENS¹ and PEYMAN KABOLIZADEH¹

¹Department of Radiation Oncology, Beaumont Health, Royal Oak, MI, U.S.A.;

²Department of Radiation Oncology, UPMC Hillman Cancer Center, Pittsburgh, PA, U.S.A.;

³Department of Chemistry, University of Pittsburgh, Pittsburgh, PA, U.S.A.

Abstract. *Background/Aim:* We tested JP4-039, a GS-nitroxide radiation damage mitigator in proton therapy of Fanconi anemia (FA) mice. *Materials and Methods:* *Fanca*^{-/-} and *Fanca*^{+/+} bone marrow stromal cells were pre-treated with JP4-039 and irradiated with either protons or photons (0-10 GyRBE) followed by clonogenic survival and β -Galactosidase senescence analysis. *Fanca*^{-/-} and *Fanca*^{+/+} mice were pretreated with JP4-039 for 10 min prior to oropharyngeal irradiation with either protons or photons (0 or 30 GyRBE) followed by sacrifice and measurement of oral cavity ulceration, distant hematopoietic suppression, and real-time polymerase chain reaction analysis. *Results:* JP4-039 reduced oral cavity ulceration in *Fanca*^{-/-} mice, transcripts *Nfkb*, *Ap1*, *Sp1*, and *Nrf2*, and proton therapy induced distant marrow suppression. *Conclusion:* JP4-039 protected *Fanca*^{-/-} and *Fanca*^{+/+} cells and mouse oral cavity from both proton and photon radiation.

Fanconi anemia (FA) is an inherited disorder that is associated with anemia, congenital abnormalities and radiosensitivity. FA patients have an increased incidence of malignancy, in particular oral cavity squamous cell carcinomas. While the incidence of FA is low, an estimated 30 patients in the USA each year (1), survival of these patients has been prolonged due to advances in hematopoietic cell transplantation. Consequently, more FA

patients are developing malignancies, particularly squamous cell carcinomas of the head, neck, esophagus, and urogenital regions (2). Some FA patients are sensitive to radiation therapy due to a mutation in one or more of the 23 genes in the FA pathway (3). Radiation protective agents and normal tissue sparing proton therapy may help treat these patients safely. We tested the combination of JP4-039, a GS-nitroxide radiation damage mitigator, and proton therapy in a mouse model of FA.

Prior studies have demonstrated that intraoral administration of mitochondrial localized transgene protein manganese superoxide dismutase (SOD2) prior to daily fractionated RT decreased both acute and late oropharyngeal toxicity without affecting tumor control (4-6). The 4-Amino-Tempo (4-AT) demonstrated radioprotection; however, was limited by the need for high systemic doses (7, 8). A Gramicidin S (GS)-derived peptide isostere conjugated to 4-Amino-Tempo (4-AT; JP4-039) targets the nitroxide to the mitochondria, and, in several FA mouse models, allows for safe radioprotection (9-12). Mice exposed to a dose that is lethal to 50% of mice within 30 days (LD50/30) of total body irradiation (TBI) were protected by intraperitoneal (IP) injection of JP4-039 after 24 h of exposure (13, 14). In addition, in the radiosensitive *Fancd2*^{-/-}, *Fanca*^{-/-} and *Fancg*^{-/-} mice, pre-treated with oral JP4-039, there was decreased oral cavity mucositis following single fraction or fractionated therapy (10, 14).

Proton therapy has been touted as a major therapeutic approach for a number of clinical scenarios, including: dose escalation, re-irradiation, and pediatric oncology. As protons traverse tissue, they slow-down and deliver the majority of their energy when they come to rest, a phenomenon referred to as the Bragg peak. Pencil beam scanning (PBS) is an advanced delivery method in which individual "pencil" beams, of varied energy, are used to "paint" dose on the desired target, layer by layer, allowing for complex treatment

This article is freely accessible online.

Correspondence to: Peyman Kabolizadeh, MD, Ph.D., Department of Radiation Oncology, 3577 West 13 Mile Road, Suite LL, Royal Oak, MI 48073, U.S.A. Tel: +1 2485525490, e-mail: peyman.kabolizadeh@beaumont.org

Key Words: Fanconi anemia, proton, radiation protector, mucositis, toxicity.

volumes, and improved integral dose and novel approaches such as intensity modulated proton therapy (IMPT) to further improve treatment delivery while minimizing normal tissue exposure (15, 16).

In the present study, we sought to confirm the radioprotective effect of JP4-039 in FA mice, both *in vitro* and *in vivo*, using PBS proton therapy. Given the significant physical differences between protons and photons, we compared proton and photon therapy with respect to biological markers, including: distant bone marrow suppression, radiation induction of cell senescence and alterations in the level of RNA transcription for stress response and inflammation-related markers. To our knowledge, this is the first study to compare biological differences between photon and protons in FA *in vitro* and *in vivo* models.

Materials and Methods

Animal care. Fanca^{-/-} mice (129/Sv) were generously provided by Dr. Markus Grompe (Oregon Health Sciences Center, Portland, OR, USA) and maintained four per cage according to Institutional Animal Use and Care Committee (IACUC) protocols and fed with a standard laboratory chow diet with deionized water. Briefly, Fanca^{-/-} were generated by mutagenesis of the Fanca gene *via* replacement of exons 4-7 by the LacZ-Neo fusion marker in E14 ES cells, derived from the 129Ola strain, resulting in embryonic stem cell clones. Heterozygotes were inbred to generate homozygous mutant Fanca^{-/-} offspring, as previously described in detail (17).

Bone marrow cultures. Bone marrow stromal cells (BMSC) were collected from the femora and tibiae of Fanca^{-/-} and control Fanca^{+/+} mice and cultured in T25 cm² plastic flasks (Corning® Inc., Corning, NY, USA) with McCoy's 5A media (Gibco, Gaithersburg, MD, USA) supplemented with 25% fetal calf serum (Cambrex, Rockland, ME, USA), 10⁻⁵ M hydrocortisone hemisuccinate, and 1% penicillin-streptomycin (Gibco). Cultures were maintained at 33°C in 7% CO₂, as described previously (8).

Clonogenic radiation survival curves. BMSCs derived from Fanca^{-/-} and control Fanca^{+/+} mice were maintained in complete media in T25 flasks and were stacked and irradiated to doses of 0-10 GyRBE *via* pencil beam scanning on an IBA Proteus® One proton therapy system (Ion Beam Applications SA, Louvain-La-Neuve, Belgium). Subgroups of cultures were pre-treated with 100 μM JP4-039 for 12 h prior to irradiation. Parallel cultures were also irradiated using a JL Shepherd Mark I Model 68A 137CS γ Irradiator (San Fernando, CA, USA) at 340 cGy per min along with subgroups treated with JP4-039 in a similar fashion.

The single-hit/multi-target model was used, and results are presented as the mean of at least three experiments, as previously described (18). The dose required to reduce the surviving fraction to 37% is defined as D0. The back extrapolation to the y-axis from the linear portion of the survival curve is defined as \bar{n} . The survival curve is a log-to-linear plot with surviving fraction along the y axis (log) and radiation dose (GyRBE) along the x axis.

β-Galactosidase senescence assay. BMSCs from Fanca^{-/-} and control Fanca^{+/+} mice were irradiated with a dose of 10 GyRBE in T25 flasks using proton or photon radiation, respectively. Cells were

plated in T25 flasks and evaluated after 5 days. β-galactosidase staining was completed as described in prior publications (18).

In vivo radiation. For proton therapy experiments, 8-10-week-old Fanca^{-/-} and control Fanca^{+/+} mice were packaged by the Division of Laboratory Animal Resources (DLAR) at UPMC and shipped from Pittsburgh, PA to Royal Oak, MI, USA *via* courier. Once arrived, the mice were given time to acclimate prior to irradiation. Mice were given 100 μl water alone or an additional 100 μl of JP4-039 (20 mg/kg), 4-AT/Miglyol (20 mg/kg), or Miglyol by oral gavage 10 min prior to irradiation. The mice were immobilized in modified 50 ml conical tubes. Once immobilized, the cast and mice were scanned in a Philips Big Bore helical CT scanner for proton planning. RayStation version 6 (RaySearch AB, Stockholm, Sweden) Treatment Planning System (TPS) was used. The wax material relative proton stopping power was used in the TPS directly. A uniform spread-out Bragg peak (SOBP) of 30 GyRBE was prescribed to the head and neck region to ensure homogenous dose coverage while sparing the remainder of the treated mice. Following RT, the mice were repackaged and shipped back to Pittsburgh. For photon therapy experiments, mice were treated with a dose of 30 GyRBE, as described previously (12).

Histopathologic assessment of oral cavity tissues. Mice treated with large doses of radiation become increasingly malnourished, due to radiation-induced oral mucositis, with associated mortality occurring at approximately 6 days post-irradiation. Moreover, the IACUC recommend that mice be euthanized if there is a >20% loss of body weight. Therefore, mice were sacrificed 5 days following irradiation.

Following sacrifice, the tongue was dissected and removed from the oropharynx and fixed in 2% PFA for 2 h and then stored in 30% sucrose. Next, blocks were sectioned in 5 μm slices and stained with hematoxylin and eosin (H&E). Slides were then scored, in a blinded fashion by two observers, for percentage ulceration, as described in prior publications (6, 12).

Abscopal hematopoietic colony-forming assays. Mice were sacrificed 5 days following head and neck irradiation and single cell suspensions of fresh bone marrow were obtained from non-irradiated tibiae and femora and plated in triplicate in 0.8% methylcellulose containing Iscove's medium supplemented with hematopoietic growth factors: recombinant murine stem cell factor, IL-3, IL-6, erythropoietin (Stemcell Technologies, Vancouver, Canada). The cells were assayed for multilineage colony-forming unit-granulocyte macrophage (CFU-GM) and burst-forming unit erythroid (BFU-e) at day 13. Colonies containing ≥50 cells were scored.

Real-time polymerase chain reaction (RT-PCR) analysis of RNA transcripts. Mice were sacrificed 5 days following head and neck irradiation and tongue tissues from nonirradiated controls, radiation alone and from pretreatment JP4-039, 4-AT, and Miglyol were prepared for RT-PCR. Tissue from each mouse was prepared as previously described (12). RT-PCR analysis was performed to measure radiation-inducible transcripts of TGF-β, p53, p21, Rad51, SP-1, Sod1, Gadd45, Nfkb, IL-1α, AP-1, and Nrf2. The results were standardized with respect to glyceraldehyde phosphate dehydrogenase (GAPDH). Moreover, the results are presented as fold increase in gene transcript expression above the baseline level, which was defined as that of nonirradiated wild-type tongue tissue.




Figure 1. Clonogenic survival curves for *Fanca*^{-/-} and *Fanca*^{+/+} BMSCs treated with either proton therapy (A) or photon therapy (B) in the presence or absence of JP4-039.

Results

In vitro clonogenic radiation sensitivity assay. Clonogenic survival curves revealed that *Fanca*^{-/-} bone marrow derived stromal cells were more radiosensitive to both proton (Figure 1A) and photon (Figure 1B) radiotherapy compared to wild-type *Fanca*^{+/+} cells. The most pronounced difference observed was the dose required to reduce clonogenic survival by 37% (D0). Specifically, *Fanca*^{-/-} cells treated with proton therapy had a D0=2.11±0.17 GyRBE; while *Fanca*^{+/+} cells had a D0=3.28±0.10 GyRBE (*p*=0.0027). The shoulder (\bar{n}) was significantly narrower in *Fanca*^{-/-} cells treated with photon radiation compared to that in *Fanca*^{+/+} cells (2.0±0.5 vs. 5.7±1.3, *p*=0.029).

Pre-treatment of bone marrow derived stromal cells with 10 μM JP4-039 for 12 h prior to irradiation provided significant radiation protection for *Fanca*^{-/-} and *Fanca*^{+/+} cell lines against both proton and photon radiation. For instance, *Fanca*^{-/-} cells that were pre-treated with JP4-039 and then underwent proton radiation had a D0=3.01±0.19 GyRBE compared to D0=2.11±0.17 GyRBE for *Fanca*^{-/-} cells treated with proton therapy alone (*p*=0.0125). The radioprotective effect of JP4-039 was observed in *Fanca*^{-/-} cells treated with photon therapy as well.

In vitro β-galactosidase senescence assay. The percent of clonogenic cells stained for β-galactosidase was significantly increased following 10 GyRBE proton or photon radiation, a finding that was statistically significant compared to no radiation therapy among both *Fanca*^{+/+} and *Fanca*^{-/-} cells (Figure 2A). Interestingly, *Fanca*^{-/-} cells that were treated with proton therapy were found to have statistically significant lower proportion of β-galactosidase positive clonogens compared to *Fanca*^{+/+} cells treated at the same radiation dose (10% vs. 28%, *p*<0.05). Conversely, *Fanca*^{-/-} cells that were treated with photon therapy had a statistically significantly higher proportion of β-galactosidase positive clonogens compared to *Fanca*^{+/+} cells treated at the same radiation dose (17% vs. 7%, *p*<0.05).


We next tested *Fanca*^{+/+} and *Fanca*^{-/-} BMSCs (Figure 2B) for senescence after different doses of proton radiation and in the presence or absence of the radiation mitigator, JP4-039. The addition of JP4-039 did not affect the percent of cells stained positive for β-galactosidase at a dose level of 0, 5 or 10 GyRBE in either genotype. Interestingly, there was a statistically significant difference between *Fanca*^{-/-} mice treated to 10 GyRBE with JP4-039 and *Fanca*^{+/+} mice treated at the same radiation dose and JP4-039 concentration (*p*<0.04).

Amelioration of radiation induced mucositis with JP4-039. *Fanca*^{-/-} and *Fanca*^{+/+} mice underwent oral cavity proton radiation therapy to 30 GyRBE and degree of mucositis was assessed 5 days later (Figure 3A). The percent oral mucositis was significantly elevated compared to no radiation in both the *Fanca*^{+/+} (23.5% vs. 0.7%, *p*<0.05) and *Fanca*^{-/-} mice (26.5% vs. 2.9%, *p*<0.05) (Figure 3B). However, the degree of oral mucositis among the *Fanca*^{-/-} mice pre-treated with JP4-039 was lower compared to the mice treated with 30 GyRBE proton therapy alone (6.5% vs. 26.5%, *p*<0.0001). Notably, pre-treatment with the vehicle (Miglyol) or non-mitochondrial targeted 4-AT/Miglyol did not provide any significant reduction in oral mucositis.

Proton induced changes in oral mucosae transcripts. Proton irradiation of the oral cavity of *Fanca*^{+/+} and *Fanca*^{-/-} mice resulted in robust and statistically significant increase in Nfkb, Ap1, Sp1, and Nrf2 transcripts, relative to unirradiated oral cavity mucosa (Figure 4A-D). This pattern was recapitulated in both mice genotypes. In addition, pre-treatment with JP4-039 resulted in a statistically significant reduction in transcript levels compared to proton alone irradiated oral mucosa in *Fanca*^{+/+} mice, but not among *Fanca*^{-/-} mice. Similarly, both cytokine transcripts (TGF-β and IL-1α) and the oxidative stress response transcript Sod2 were robustly increased in proton irradiated oral mucosa of both *Fanca*^{+/+} and *Fanca*^{-/-} mice. Pre-treatment with JP4-039 reduced TGF-β and Sod1 transcript levels compared to 30 GyRBE alone in *Fanca*^{+/+} mice but not in *Fanca*^{-/-} mice. In addition, pre-treatment with JP4-039 reduced IL-1α transcript levels in *Fanca*^{-/-} mice but not in *Fanca*^{+/+} mice.


Finally, the radiation response-related transcripts Gadd45, Rad51, p21 and p53 were increased following proton irradiation of *Fanca*^{+/+} and *Fanca*^{-/-} oral mucosa, except p53 in *Fanca*^{+/+} mice (Figure 5A-D). Moreover, pre-treatment with JP4-039 did not significantly reduce the levels of these transcripts compared to levels seen in proton irradiated oral mucosa, except for Gadd45 in *Fanca*^{-/-} mice. Of note, JP4-039 modulated Rad51 and p53 transcript levels to a level similar to that of the nonirradiated oral mucosa of *Fanca*^{+/+} mice.

Radiation induced bone marrow suppression and mitigation of the suppression by JP4-039. *Fanca*^{+/+} and *Fanca*^{-/-} mice that received 30 GyRBE to the oral cavity developed distal bone marrow suppression when evaluated with CFU-GM (Figure 6A) and BFU-e (Figure 6B) colony forming assays. The absolute difference in CFU-GM cells between no radiation and 30 GyRBE was similar for both *Fanca*^{+/+} and *Fanca*^{-/-} mice. In contrast, the magnitude of distal marrow suppression, reflected by BFU-e cells, was much higher in the *Fanca*^{-/-} mice, while the difference was not statistically significant in *Fanca*^{+/+} mice.



* *p* < 0.05 compared to 0 GyRBE of the same genotype
p < 0.05 compared to *Fanca* +/+ of the same irradiation dose

□ 0 GyRBE
▨ 10 GyRBE



* *p* < 0.05 compared to 0 GyRBE of the same genotype/drug concentration

p < 0.04 compared to *Fanca* +/+ and same radiation dose/drug concentration

■ 0mM JP4-039
■ 10mM JP4-039

Figure 2. β-Galactosidase assay for *Fanca*^{+/+} or *Fanca*^{-/-} BMSCs following (A) proton or photon radiation therapy. Effect of (B) JP4-039 on proton-induced senescence in both FA models.




Figure 3. *Fanca*^{+/+} and *Fanca*^{-/-} mice were irradiated with 30 GyRBE to the oral cavity, in the presence or absence of vehicle, non-targeted 4-AT or JP4-039, and the percent of mucositis was assessed (A/B).

Pre-treatment with JP4-039 reduced proton-induced distal marrow suppression in both mice models. Moreover, there was a statistically significant increase in CFU-GM cells compared to the all subgroups for both *Fanca*^{+/+} and *Fanca*^{-/-} mice ($p < 0.05$). There was a reduction in CFU-GM cells in response to 30 GyRBE relative to non-irradiated

controls, indicative of abscopal bone marrow suppression. Pre-treatment of proton-treated *Fanca*^{+/+} (Figure 6C) and *Fanca*^{-/-} (Figure 6D) mice with JP4-039 increased marrow recovery relatively to proton therapy alone ($p < 0.0001$) and compared to the photon treatment + JP4-039 group ($p = 0.0014$).




Figure 4. Proton-induced changes in the pro-inflammatory transcription factor transcripts (A) *Nfkb*, (B) *Ap1*, (C) *Sp1* and (D) *Nrf2* are mitigated by oral administration of JP4-039 in both FA models.

Discussion

FA patients are at high risk of developing squamous cell carcinoma of the head and neck. With the advent of stem cell transplantation and the increased life expectancy more FA patients are at risk. FA patients with locally advanced oral cavity cancer often require a combination of surgery, radiation therapy and chemotherapy. There are subsets of FA patients that are highly radiosensitive, and there is no established biomarker to identify these patients. Therefore, novel interventions, which spare and/or protect normal tissue are needed. To that end, we have extended the preclinical efficacy of JP4-039 in photon irradiation (10-12) by evaluating its utility in *Fanca*^{-/-} and *Fanca*^{+/+} mice models irradiated with a high single fraction of proton radiation.

The International Commission on Radiation Units (ICRU)-78 report assumes a relative biological effectiveness (RBE) of 1.1 for protons, while heavier particles are reported to have a variable RBE (19). However, a fixed RBE for protons is based on assumptions of macroscopic and homogeneous dose

in a volume of interest, which do not remain valid at the microscopic level (20). Therefore, the biological consequences of proton therapy are possibly different at the microscopic level when compared to photon radiation (21). In the current study, we observed photon curves demonstrating a characteristic shoulder while the proton curve had a more linear shape (Figure 1A and B). These differences may reflect an inherent difference in the biological response to proton therapy.

Recent publications suggest that senescence is one of multiple cellular responses to ionizing irradiation both *in vitro* and *in vivo*. *Fanca*^{-/-} and *Fanca*^{+/+} stromal cell lines were irradiated *in vitro* and serial cultures were then tested for senescence by β -galactosidase staining according to published methods (22). As shown in Figure 2, both *Fanca*^{-/-} and *Fanca*^{+/+} BMSCs demonstrated proton- and photon-induced increase in the percent of senescent cells quantitated by β -galactosidase staining 5 days post-irradiation. However, the relative increase in proton irradiated *Fanca*^{-/-} cells was lower compared to photon irradiated *Fanca*^{-/-} cells. The data suggest




Figure 5. Proton-induced changes in the radiation response-related transcripts (A) *Gadd45*, (B) *Rad51*, (C) *p21*, and (D) *p53* in both FA models.

putative biological differences in the response of cells to photon compared to proton irradiation.

While β -galactosidase is one of multiple markers of senescence, other studies measuring levels of p16, p21 and telomere length have also been used as biomarkers for quantitating senescence. Future studies will determine whether these three biomarkers follow the same pattern as that observed with β -galactosidase and may confirm the biological difference in the response between proton and photon irradiated *Fanca*^{-/-} and *Fanca*^{+/+} cell lines. Moreover, fractionated irradiation dose response curves may further elucidate potential differences in the molecular mechanisms of senescence pathways between protons and photons.

We have published extensively on the radioprotective effects of JP4-039 in cell culture and mouse models (12, 14, 19, 23-26). In the present report, we demonstrated that orally administered JP4-039, prior to head and neck proton irradiation, mitigates radiation-induced tongue ulceration in both *Fanca*^{+/+} and *Fanca*^{-/-} models. At the genetic level, pretreatment with JP4-039 differentially modulated transcription levels, with reduction in TGF- β , Sod1, Nfkb, Ap1, Sp1 and

Nrf2 among *Fanca*^{+/+} mice and IL-1 α among *Fanca*^{-/-} mice. In contrast, pre-treatment with JP4-039 did not significantly reduce the transcript levels in *Fanca*^{-/-} mice. The magnitude of changes in proton irradiated oral mucosa was lower compared to photon radiation. These results further extend that of prior publications demonstrating a therapeutic effect of JP4-039 with respect to reduction of irradiation-induced transcripts of stress response genes and inflammatory cytokines (10-12).

We have previously demonstrated that single-fraction head and neck photon radiation results in distal bone marrow suppression in both *Fandc2*^{+/+} and *Fandc2*^{-/-} mice, and this is reduced with JP4-039 pretreatment (10-12). The current study further extended these findings and demonstrated that JP4-039 reduces proton induced distant bone marrow suppression as well. Additionally, JP4-039 with proton therapy resulted in a statistically significant relative increase in bone marrow activity when compared to JP4-039 plus photon radiation, supporting other possible biological differences between photon and proton irradiation.

JP4-039 was able to protect *Fanca*^{-/-} and *Fanca*^{+/+} cells from both proton and photon radiation. This was also seen in




Figure 6. Distant bone marrow suppression [(A) CFU-GM and (B) BFU-e] following oral cavity proton irradiation of *Fanca*^{+/+} and *Fanca*^{-/-} mice, relative to no radiation (C-D) in the presence or absence of JP4-039.

our *in vivo* model, in which oral JP4-039 reduced the degree of oral mucositis following a large single fraction of 30 GyRBE to the head and neck region. These data support future clinical trials to evaluate the use of intraoral JP4-049 among FA patients receiving radiation therapy and indicate that more research is required to understand the differences between photon and proton irradiation.

Conflicts of Interest

Dr. Greenberger reports grants from NIH, during the conduct of the study; Dr. Wipf reports grants from NIH, during the conduct of the study; In addition, Dr. Wipf has a patent WO 2012068081 A1 20120524 issued; Dr. Epperly reports grants from NIH, during the conduct of the study. There are no other conflicts of interest to disclose.

Authors' Contribution

Conception and design: TJ Quinn, GD Wilson, MW Epperly, JS Greenberger, CW Stevens, P Kabolizadeh; Administrative support: X Ding, GD Wilson, MW Epperly, D Franciola, JS Greenberger, CW Stevens, P Kabolizadeh; Provision of study materials or patients: X Ding, X Li, GD Wilson, MW Epperly, P Wipf, JS Greenberger, CW Stevens, P Kabolizadeh; Collection and assembly

of data: TJ Quinn, K Buelow, A Sivananthan, S Thermoziar, A Henderson, M Epperly, JS Greenberger, CW Stevens, P Kabolizadeh; Data analysis and interpretation: TJ Quinn, GD Wilson, MW Epperly, P Wipf, JS Greenberger, CW Stevens, P Kabolizadeh; Manuscript writing: TJ Quinn, GD Wilson, MW Epperly, P Wipf, JS Greenberger, CW Stevens, P Kabolizadeh; Final approval of manuscript: TJ Quinn, X Ding, X Li, GD Wilson, K Buelow, A Sivananthan, S Thermoziar, A Henderson, MW Epperly, D Franciola, P Wipf, JS Greenberger, CW Stevens, P Kabolizadeh.

Acknowledgements

This work was funded by the NIH grant U19-A168021.

References

- Collins J and Dokal I: Inherited bone marrow failure syndromes. *Hematology (Amsterdam, Netherlands)* 20(7): 433-434, 2015. PMID: 26218202. DOI: 10.1179/1024533215Z.000000000381
- Kutler DI, Singh B, Satagopan J, Batish SD, Berwick M, Giampietro PF, Hanenberg H and Auerbach AD: A 20-year perspective on the international fanconi anemia registry (IFAR). *Blood* 101(4): 1249-1256, 2003. PMID: 12393516. DOI: 10.1182/blood-2002-07-2170

- 3 Birkeland AC, Auerbach AD, Sanborn E, Parashar B, Kuhel WI, Chandrasekharappa SC, Smogorzewska A and Kutler DI: Postoperative clinical radiosensitivity in patients with fanconi anemia and head and neck squamous cell carcinoma. *Arch Otolaryngol Head Neck Surg* 137(9): 930-934, 2011. PMID: 21930984. DOI: 10.1001/archoto.2011.154
- 4 Dhanasekaran A, Kotamraju S, Karunakaran C, Kalivendi SV, Thomas S, Joseph J and Kalyanaraman B: Mitochondria superoxide dismutase mimetic inhibits peroxide-induced oxidative damage and apoptosis: Role of mitochondrial superoxide. *Free Radic Biol Med* 39(5): 567-583, 2005. PMID: 16085176. DOI: 10.1016/j.freeradbiomed.2005.04.016
- 5 Epperly MW, Wegner R, Kanai AJ, Kagan V, Greenberger EE, Nie S and Greenberger JS: Effects of MnSOD-plasmid liposome gene therapy on antioxidant levels in irradiated murine oral cavity orthotopic tumors. *Radiat Res* 167(3): 289-297, 2007. PMID: 17316075. DOI: 10.1667/RR0761.1
- 6 Guo H, Seixas-Silva JA, Epperly MW, Gretton JE, Shin DM, Bar-Sagi D, Archer H and Greenberger JS: Prevention of radiation-induced oral cavity mucositis by plasmid/liposome delivery of the human manganese superoxide dismutase (SOD2) transgene. *Radiat Res* 159(3): 361-370, 2003. PMID: 12600239. DOI: 10.1667/0033-7587(2003)159[0361:PORIOC]2.0.CO;2
- 7 Xavier S, Yamada K, Samuni AM, DeGraff W, Krishna MC and Mitchell JB: Differential protection by nitroxides and hydroxylamines to radiation-induced and metal ion-catalyzed oxidative damage. *Biochim Biophys Acta* 1573(2): 109-120, 2002. PMID: 12399020. DOI: 10.1016/s0304-4165(02)00339-2
- 8 Hahn SM, Krishna MC, DeLuca AM, Coffin D and Mitchell JB: Evaluation of the hydroxylamine tempol-H as an *in vivo* radioprotector. *Free Radic Biol Med* 28(6): 953-958, 2000. PMID: 10802227. DOI: 10.1016/s0891-5849(00)00176-3
- 9 Jiang J, Belikova NA, Hoye AT, Zhao Q, Epperly MW, Greenberger JS, Wipf P and Kagan VE: A mitochondria-targeted nitroxide/hemigramicidin S conjugate protects mouse embryonic cells against gamma irradiation. *Int J Radiat Oncol Biol Phys* 70(3): 816-825, 2007. PMID: 18262096. DOI: 10.1016/j.ijrobp.2007.10.047
- 10 Berhane H, Shinde A, Kalash R, Xu K, Epperly MW, Goff J, Francicola D, Zhang X, Dixon T, Shields D, Wang H, Wipf P, Li S, Gao X and Greenberger JS: Amelioration of radiation-induced oral cavity mucositis and distant bone marrow suppression in fanconi anemia *Fancd2*^{-/-} (FVB/N) mice by intraoral GS-nitroxide JP4-039. *Radiat Res* 182(1): 35-49, 2014. PMID: 24932534. DOI: 10.1667/RR13633.1
- 11 Willis J, Epperly MW, Fisher R, Zhang X, Shields D, Hou W, Wang H, Li S, Wipf P, Parmar K, Guinan E, Steinman J and Greenberger JS: Amelioration of head and neck radiation-induced mucositis and distant marrow suppression in *Fanca*^{-/-} and *fancg*^{-/-} mice by intraoral administration of GS-nitroxide (JP4-039). *Radiat Res* 189(6): 560-578, 2018. PMID: 29584588. DOI: 10.1667/RR14878.1
- 12 Shinde A, Berhane H, Rhieu BH, Kalash R, Xu K, Goff J, Epperly MW, Francicola D, Zhang X, Dixon T, Shields D, Wang H, Wipf P, Parmar K, Guinan E, Kagan V, Tyurin V, Ferris RL, Zhang X, Li S and Greenberger JS: Intraoral mitochondrial-targeted GS-nitroxide, JP4-039, radioprotects normal tissue in tumor-bearing radiosensitive *Fancd2*^(-/-) (C57BL/6) mice. *Radiat Res* 185(2): 134-150, 2016. PMID: 26789701. DOI: 10.1667/RR14035.1
- 13 Goff JP, Epperly MW, Dixon T, Wang H, Francicola D, Shields D, Wipf P, Li S, Gao X and Greenberger JS: Radiobiologic effects of GS-nitroxide (JP4-039) on the hematopoietic syndrome. *In Vivo* 25(3): 315, 2011. PMID: 21576404.
- 14 Epperly MW, Wipf P, Fisher R, Francicola D, Beumer J, Li S, Brand RM, Falo LD Jr, Erdos G and Greenberger JS: Evaluation of different formulations and routes for the delivery of the ionizing radiation mitigator GS-nitroxide (JP4-039). *In Vivo* 32(5): 1009-1023, 2018. PMID: 30150422. DOI: 10.21873/invivo.11341
- 15 Ding X, Li X, Zhang JM, Kabolizadeh P, Stevens C and Yan D: Spot-scanning proton arc (SPArc) therapy: The first robust and delivery-efficient spot-scanning proton arc therapy. *Int J Radiat Oncol Biol Phys* 96(5): 1107-1116, 2016. PMID: 27869083. DOI: 10.1016/j.ijrobp.2016.08.049
- 16 Li X, Kabolizadeh P, Yan D, Qin A, Zhou J, Hong Y, Guerrero T, Grills I, Stevens C and Ding X: Improve dosimetric outcome in stage III non-small-cell lung cancer treatment using spot-scanning proton arc (SPArc) therapy. *Radiat Oncol* 13(1): 35, 2018. PMID: 29486782. DOI: 10.1186/s13014-018-0981-6
- 17 Cheng NC, van de Vrugt HJ, van der Valk MA, Oostra AB, Krimpenfort P, de Vries Y, Joenje H, Berns A and Arwert F: Mice with a targeted disruption of the fanconi anemia homolog *fanca*. *Hum Mol Genet* 9(12): 1805-1811, 2000. PMID: 10915769. DOI: 10.1093/hmg/9.12.1805
- 18 Rwigema JM, Beck B, Wang W, Doemling A, Epperly MW, Shields D, Francicola D, Dixon T, Frantz M-C, Wipf P, Tyurina Y, Kagan VE, Wang H and Greenberger JS: Two strategies for the development of mitochondrial-targeted small molecule radiation damage mitigators. *Int J Radiat Oncol Biol Phys* 80(3): 860-868, 2011. PMID: 21493014. DOI: 10.1016/j.ijrobp.2011.01.059
- 19 International Commission on Radiation Units and Measurements: ICRU Report 78. *J ICRU* 7(2), 2007. Oxford University Press, Oxford, 2007.
- 20 Paganetti H: Relative biological effectiveness (RBE) values for proton beam therapy. variations as a function of biological endpoint, dose, and linear energy transfer. *Phys Med Biol* 59(22): R472, 2014. PMID: 25361443. DOI: 10.1088/0031-9155/59/22/R419
- 21 van Luijk P and Paganetti H: Biological considerations when comparing proton therapy with photon therapy. *Semin Radiat Oncol* 23(2): 77-87, 2013. PMID: 23473684. DOI: 10.1016/j.semradonc.2012.11.002
- 22 Sivananthan A, Shields D, Fisher R, Hou W, Zhang X, Francicola D, Epperly MW, Wipf P and Greenberger JS: Continuous one-year oral administration of the radiation mitigator, MMS350, after total-body irradiation, restores bone marrow stromal cell proliferative capacity and reduces senescence in fanconi anemia (*Fanca*^{-/-}) mice. *Radiat Res* 191(2): 139-153, 2019. PMID: 30499383. DOI: 10.1667/RR15199.1
- 23 Epperly MW, Goff JP, Li S, Gao X, Wipf P, Dixon T, Wang H, Francicola D, Shen H, Rwigema JC, Kagan V, Bernard M and Greenberger JS: Intraesophageal administration of GS-nitroxide (JP4-039) protects against ionizing irradiation-induced esophagitis. *In Vivo* 24(6): 811-819, 2010. PMID: 21164038.
- 24 Bernard ME, Kim H, Berhane H, Epperly MW, Francicola D, Zhang X, Houghton F, Shields D, Wang H, Bakkenist CJ, Frantz MC, Forbeck EM, Goff JP, Wipf P and Greenberger JS: GS-nitroxide (JP4-039)-mediated radioprotection of human fanconi anemia cell lines. *Radiat Res* 176(5): 603-612, 2011. PMID: 21939290. DOI: 10.1667/RR2624.1

- 25 Epperly MW, Goff JP, Franicola D, Wang H, Wipf P, Li S and Greenberger JS: Esophageal radioprotection by swallowed JP4-039/F15 in thoracic-irradiated mice with transgenic lung tumors. *In Vivo* 28(4): 435-440, 2014. PMID: 24982207.
- 26 Hebish Berhane, Michael W. Epperly, Julie Goff, Kalash R, Cao S, Franicola D, Zhang X, Shields D, Houghton F, Wang H, Wipf P, Parmar K and Greenberger JS: Radiologic differences between

bone marrow stromal and hematopoietic progenitor cell lines from fanconi anemia (*Fancd2*^{-/-}) mice. *Radiation Res* 181(1): 76-89, 2014. PMID: 24397476. DOI: 10.1667/RR13405.1

Received August 13, 2019

Revised September 14, 2019

Accepted September 16, 2019