

Editorial Board

- I. ABRAHAM, *The HOPE Center, University of Arizona, Tucson, AZ, USA*
- N.J. AGNANTIS, *Department of Pathology, University of Ioannina, Ioannina, Greece*
- D. ANDERSON, *Department of Biomedical Sciences, University of Bradford, Bradford, West Yorkshire, UK*
- V. BARAK, *Department of Oncology, Hadassah University Hospital, Jerusalem, Israel*
- M.H. BARCELLOS-HOFF, *Department of Radiation Oncology, New York University School of Medicine, New York, NY, USA*
- K. BEIER, *Department of Histology, University of Basel, Basel, Switzerland*
- M. BERGQVIST, *Department of Oncology, Radiology and Clinical Immunology, University Hospital, Uppsala, Sweden*
- R. BJERKVIG, *Norlux Neuro-Oncology, Department of Biomedicine, University of Bergen, Bergen, Norway*
- W. BOWNE, *Department of Surgery, Drexel University College of Medicine, Philadelphia, PA, USA*
- M. CARAGLIA, *Department of Experimental Oncology, National Institute of Tumours Fondazione G. Pascale, Naples, Italy*
- P. CHANDRA, *Department of Molecular Biology, Frankfurt University, Frankfurt am Main, Germany*
- J.-G. CHUNG, *Department of Medicine, China Medical College, Taichung, Taiwan, ROC*
- L.A. COHEN, *Northampton, MA, USA*
- A.I. CONSTANTINOU, *Department of Biological Sciences, University of Cyprus, Nicosia, Cyprus*
- T. DALIANIS, *Department of Pathology-Oncology, Karolinska Institute, Stockholm, Sweden*
- D.T. DENHARDT, *Division of Life Sciences, Rutgers University, Bridgewater, NJ, USA*
- W. DEN OTTER, *VUMC - Department of Urology, Amsterdam, The Netherlands*
- K. DE MEIRLEIR, *Department of Human Physiology and Medicine, Faculty of Physical Education and Physiotherapy, Vrije Universiteit Brussel, Brussels, Belgium*
- L. DE RIDDER, *Department of Anatomy, Embryology and Histology, University of Ghent, Ghent, Belgium*
- E.P. DIAMANDIS, *Department of Pathology and Laboratory Medicine, University of Toronto, Toronto, Ontario, Canada*
- J.M. DRAKE, *Division of Medical Oncology, RBHS-Robert Wood Johnson Medical School, Rutgers Cancer Institute of New Jersey, New Brunswick, NJ, USA*
- T. EFFERTH, *Department of Pharmaceutical Biology, Institute of Pharmacy and Biochemistry, University of Mainz, Germany*
- W. ENGSTRÖM, *Department of Molecular Biosciences, Swedish University of Agricultural Sciences, Uppsala, Sweden*
- M. ESKELINEN, *Department of Surgery, University Hospital of Kuopio, Kuopio, Finland*
- J.A. FERNANDEZ-POL, *Metalloproteomics, LLC, Chesterfield, MO, USA*
- G. FIORENTINI, *UOC Oncologia, Azienda Ospedaliera Marche Nord, Pesaro, Italy*
- P.B. FISHER, *Virginia Commonwealth University, School of Medicine, Richmond, VA, USA*
- I. FREITAS, *Dipartimento di Biologia Animale, University of Pavia, Pavia, Italy*
- M. FRIEDRICH, *Department of Obstetrics and Gynecology, Klinikum Krefeld, Krefeld, Germany*
- R.E. FRIEDRICH, *Department of Oral and Maxillofacial Surgery, Eppendorf University Hospital, Hamburg, Germany*
- R. GANAPATHI, *Levine Cancer Institute, Carolinas HealthCare System, Charlotte, NC, USA*
- G. GITSCH, *Department of Gynecology and Obstetrics, University of Freiburg Medical Center, Freiburg, Germany*
- V. GORGOLIS, *Department of Histology-Embryology, Medical School, University of Athens, Greece*
- J.S. GREENBERGER, *Department of Radiation Oncology, University of Pittsburgh, Pittsburgh, PA, USA*
- J.W. GREINER, *National Cancer Institute, NIH, Bethesda, MD, USA*
- D.S. GRIDLEY, *Department of Radiation Medicine, Radiation Research Laboratories, Loma Linda University and Medical Center, Loma Linda, CA, USA*
- C.J. GRUBBS, *Department of Nutrition Sciences, University of Alabama, Birmingham, AL, USA*
- F. GUADAGNI, *IRCCS San Raffaele, Rome, Italy*

- J. HAU**, *Department of Comparative Medicine, University of Copenhagen, Denmark*
- M. HAUER-JENSEN**, *Arkansas Cancer Research Center, University of Arkansas Medical Sciences, Little Rock, AR, USA*
- K. HIBI**, *Tokai Central Hospital, Kakamigahara, Gifu, Japan*
- S.A. IMAM**, *Huntington Medical Research Institutes, Gene Therapy Program, Pasadena, CA, USA*
- K.S. JEONG**, *Department of Pathology, College of Veterinary Medicine, Kyungpook National University, Daegu, S. Korea*
- M. JHANWAR-UNIYAL**, *Department of Neurosurgery, New York Medical College, Valhalla, NY, USA*
- I. KISS**, *Institute of Preventive Medicine, Medical School, University of Pécs, Pécs, Hungary*
- E. KONDO**, *Division of Molecular and Cellular Pathology Niigata University Graduate School of Medical and Dental Sciences, Chuo-ku, Niigata, Japan*
- M. KOUTSILIERIS**, *Department of Experimental Physiology, University of Athens Medical School, Athens, Greece*
- G.R.F. KRUEGER**, *Department of Anatomy II, Center for Anatomy, The University of Cologne Medical School, Cologne, Germany*
- B. KRUSLIN**, *Department of Pathology, Ljudevit Jurak University, Sestre milosrdnice University Hospital, Zagreb, Croatia*
- N. KYPRIANOU**, *Department of Urology, University of Kentucky, Lexington, KY, USA*
- G. LANDBERG**, *Department of Laboratory Medicine, Lund University, Lund, Sweden*
- I. LELONG-REBEL**, *Lab. de Biophotonique et Pharmacologie, Université de Strasbourg, Illkirch, France*
- J. LEROY**, *IRCAD, University of Strasbourg, France*
- W. LICHTENEGGER**, *Charité Campus Virchow-Klinikum, Berlin, Germany*
- J. LYKKESFELDT**, *Department of Veterinary Disease Biology, University of Copenhagen, Frederiksberg, Denmark*
- P. MADARNAS**, *Department of Pathology, University of Sherbrooke, Québec, Canada*
- H. MAEDA**, *BioDynamics Research Foundation, Kumamoto, Japan*
- M. MAREEL**, *Laboratory of Experimental Cancerology, State University, Ghent, Belgium*
- L. MARGOLIS**, *Eunice Kennedy Shriver National Institute of Child Health and Human Development, Bethesda, MD, USA*
- G. MARTORANA**, *Policlinico S. Orsola Malpighi, University of Bologna, Bologna, Italy*
- J. MOLNÁR**, *Department of Medical Microbiology and Immunobiology, Faculty of Medicine, University of Szeged, Szeged, Hungary*
- S.L. MOOBERRY**, *Cancer Therapy & Research Center, University of Texas Health Science Center at San Antonio, San Antonio, TX, USA*
- N. MOTOHASHI**, *Department of Medicinal Chemistry, Meiji Pharmaceutical University, Tokyo, Japan*
- R.M. NAGLER**, *Department of Oral and Maxillofacial Surgery, Rambam Medical Center, Haifa, Israel*
- S. NAKANO**, *Graduate School of Health and Nutritional Sciences, Nakamura Gakuen University, Fukuoka, Japan*
- M. NAKASHIMA**, *Department of Tumor and Diagnostic Pathology, Atomic Bomb Disease Institute, Nagasaki University, Nagasaki, Japan*
- R. NARAYANAN**, *Department of Biological Sciences, Florida Atlantic University, Boca Raton, FL, USA*
- M.B. NICHOLL**, *South Texas Veterans Health Care System, San Antonio, TX, USA*
- K. NILSSON**, *Tumor Biology Laboratory, Department of Pathology, University Hospital, Uppsala, Sweden*
- K.R. NORUM**, *Institute for Nutrition Research, University of Oslo, Oslo, Norway*
- K. OGAWA**, *Department of Surgery, Tokyo Women's Medical University Medical Center East, Tokyo, Japan*
- M. PAGÉ**, *Département de Biologie, Division de Biochimie, Université Laval, Québec, Canada*
- C. POLYCHRONAKOS**, *Department of Paediatrics and Human Genetics, McGill University Health Centre Research Institute, Montreal, QC, Canada*
- M.-F. POUPON**, *Institut Curie, Section de Recherche, Paris, France*
- D. RADES**, *Department of Radiation Oncology, University of Lübeck, Lübeck, Germany*
- F.M. ROBERTSON**, *Virginia Commonwealth University, Clinical Research Services, Center for Clinical and Translational Research, Richmond Academy of Medicine, Richmond, VA, USA*

continued

- D. RUBELLO, *Department of Nuclear Medicine, PET Unit, S. Maria della Misericordia Hospital, IOV, Rovigo, Italy*
- C.A. RUBIO, *Karolinska Institute, Pathology Research Laboratory, Stockholm, Sweden*
- G.R. RUTTEMAN, *Department of Clinical Sciences of Companion Animals, University of Utrecht, Utrecht, The Netherlands*
- H. SAKAGAMI, *Department of Dental Pharmacology, Meikai University School of Dentistry, Saitama, Japan*
- G. SAVA, *Institutes of Biological Research, Fondazione Callerio-Onlus, Trieste, Italy*
- D. SCHIFFER, *Neuro-bio-oncology Center, Policlinico di Monza Foundation, University of Turin, Vercelli, Italy*
- L.D. SHULTZ, *The Jackson Laboratory, Bar Harbor, ME, USA*
- G. SICA, *Istituto di Istologia ed Embriologia, Universita Cattolica del Sacro Cuore, Roma, Italy*
- J.M. SIEGFRIED, *Department of Pharmacology, MNPI Center, University of Minnesota, Minneapolis, MN, USA*
- J. SLANSKY, *Integrated Department of Immunology, School of Medicine, University of Colorado Denver and National Jewish Health, Denver, CO, USA*
- R.M. SNAPKA, *Department of Radiology, The Ohio State University, Columbus, OH, USA*
- G.-I. SOMA, *Department of Integrated and Holistic Immunology, Faculty of Medicine, Kagawa University, Kagawa, Japan*
- P.P. SORDILLO, *Department of Medical Oncology and Hematology, Lenox Hill Hospital, New York, NY, USA*
- T.A. SPRINGER, *Department of Pathology, Harvard Medical School, Boston, MA, USA*
- D.D. SPYROPOULOS, *Department of Pathology and Laboratory Medicine, Medical University of South Carolina, Charleston, SC, USA*
- K. SYRJÄNEN, *Biohit Healthcare Oyj, Helsinki, Finland*
- G.C. TORRE, *Centro Ippocrate, Vico Cosmelli 4, Finale Ligure (SV), Italy*
- B. TRIBUKAIT, *Department of Medical Radiobiology, Karolinska Hospital, Stockholm, Sweden*
- J. VADGAMA, *Department of Internal Medicine, Charles E. Drew University of Medicine and Science, Los Angeles, CA, USA*
- J.K. VISHWANATHA, *Graduate School of Biomedical Sciences, University of North Texas Health Science Center, Fort Worth, TX, USA*
- W. WANG, *Research Institute of Healthcare Science, University of Wolverhampton, Wolverhampton, UK*
- N. WATANABE, *Department of Clinical Laboratory Medicine, Sapporo Medical University School of Medicine, Sapporo, Japan*
- W. WEBER, *Clinical Cancer Etiology Unit, Basel, Switzerland*
- L.M. WEINER, *Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, DC, USA*
- J.A. WERNER, *Department of Otolaryngology, Head and Neck Surgery, Marburg, Germany*
- S. YLÄ-HERTTUALA, *Department of Molecular Medicine, A.I. Virtanen Institute, University of Kuopio, Finland*
- H. YOSHIDA, *Department of Pathology, Kagoshima University, Kagoshima, Japan*